arXiv:1603.00932v1 [math.GN] 3 Mar 2016

A Generalization of the Stone Duality Theorem

G. Dimov, E. Ivanova-Dimova and D. Vakarelov

Department of Mathematics and Informatics, University of Sofia,

5 J. Bourchier Blvd., 1164 Sofia, Bulgaria

Abstract

We prove a new duality theorem for the category of precontact algebras which implies the Stone Duality Theorem, its connected version obtained in [16], the recent duality theorems from [3, 23], and some new duality theorems for the category of contact algebras and for the category of complete contact algebras.

1 Introduction

This paper is a continuation of the papers [13, 15, 16] and, to some extent, of the papers [7, 8, 9, 10, 11, 12, 14, 29]. In it we prove a new duality theorem for the category of precontact algebras which implies the Stone Duality Theorem, its connected version obtained in [16], the recent duality theorems from [3, 23], and some new duality theorems for the category of contact algebras and for the category of complete contact algebras. More precisely, we show that there exists a duality functor G^a between the category **PCA** of all precontact algebras and suitable morphisms between them and the category **PCS** of all 2-precontact spaces and suitable morphisms between them. Then, clearly, fixing some full subcategory $\mathcal{C}$ of the category **PCA**, we obtain a duality between the categories $\mathcal{C}$ and $G^a(\mathcal{C})$. Finding categories which are isomorphic or equivalent to the category $\mathcal{C}$ and (or) to the category $G^a(\mathcal{C})$, we obtain as corollaries the Stone Duality and the other dualities mentioned above. For example, when $\mathcal{C}$ is

[1] *Keywords:* (pre)contact algebra, 2-(pre)contact space, Stone space, Stone 2-space, (Stone) duality, C-semiregular spaces, (complete) Boolean algebra, Stone adjacency space, (closed) relations, u-points, mereocompact space.

[2] *2010 Mathematics Subject Classification:* 54E05, 18A40, 54H10, 06E15, 03G05, 54D30, 54D10.

[3] *E-mail addresses:* gdimov@fmi.uni-sofia.bg, elza@fmi.uni-sofia.bg, dvak@fmi.uni-sofia.bg

the full subcategory of the category **PCA** having as objects all contact algebras of the form (B, ρ_s^B), where $a\rho_s^B b \iff a.b \neq 0$, we obtain that the category $\mathcal{C}$ is isomorphic to the category **Bool** of Boolean algebras and Boolean homomorphisms and the category $G^a(\mathcal{C})$ is isomorphic to the category **Stone** of compact zero-dimensional Hausdorff spaces and continuous maps; in this way we obtain the Stone Duality Theorem; when $\mathcal{C}$ is the full subcategory of the category **PCA** having as objects all contact algebras of the form (B, ρ_l^B), where $a\rho_l^B b \iff a, b \neq 0$, we obtain in a similar way the connected version of the Stone Duality Theorem, described in [16].

The structure of the paper is the following. In Section 2 we give the necessary preliminary results and definitions. In Section 3 we prove our main theorem, namely, the duality theorem for precontact algebras, and obtain as corollaries the theorems mentioned above.

We now fix the notations.

All lattices are with top (= unit) and bottom (= zero) elements, denoted respectively by 1 and 0. We do not require the elements 0 and 1 to be distinct.

If $(X, \mathcal{T})$ is a topological space and M is a subset of X, we denote by $\mathrm{cl}_{(X,\mathcal{T})}(M)$ (or simply by $\mathrm{cl}(M)$ or $\mathrm{cl}_X(M)$) the closure of M in $(X, \mathcal{T})$ and by $\mathrm{int}_{(X,\mathcal{T})}(M)$ (or briefly by $\mathrm{int}(M)$ or $\mathrm{int}_X(M)$) the interior of M in $(X, \mathcal{T})$. The compact spaces are not assumed to be Hausdorff (as it is adopted in [20]).

If X is a topological space, we denote by $CO(X)$ the set of all clopen (=closed and open) subsets of X. Obviously, $(CO(X), \cup, \cap, \setminus, \emptyset, X)$ is a Boolean algebra.

If X is a set, we denote by 2^X the power set of X.

The set of all ultrafilters of a Boolean algebra B is denoted by $Ult(B)$.

If $\mathcal{C}$ denotes a category, we write $X \in |\mathcal{C}|$ if X is an object of $\mathcal{C}$, and $f \in \mathcal{C}(X, Y)$ if f is a morphism of $\mathcal{C}$ with domain X and codomain Y.

The main reference books for all notions which are not defined here are [1, 20, 24].

2 Preliminaries

We will first recall the notions of *precontact algebra* and *contact algebra*. They can be regarded as algebraic analogues of proximity spaces (see [19, 26, 4, 2, 25] for proximity spaces).

Definition 2.1. An algebraic system $\underline{B} = (B, C)$ is called a *precontact algebra* ([17]) (abbreviated as PCA) if the following holds:

- $B = (B, 0, 1, +, ., *)$ is a Boolean algebra (where the complement is denoted by "$*$");
- C is a binary relation on B (called a *precontact relation*) satisfying the following axioms:

($C0$) If aCb then $a \neq 0$ and $b \neq 0$;

($C+$) $aC(b + c)$ iff aCb or aCc; $(a + b)Cc$ iff aCc or bCc.

A precontact algebra (B, C) is said to be *complete* if the Boolean algebra B is complete. Two precontact algebras $\underline{B} = (B, C)$ and $\underline{B_1} = (B_1, C_1)$ are said to be *PCA-isomorphic* (or, simply, *isomorphic*) if there exists a *PCA-isomorphism* between them, i.e., a Boolean isomorphism $\varphi : B \longrightarrow B_1$ such that, for every $a, b \in B$, aCb iff $\varphi(a)C_1\varphi(b)$.

The negation of the relation C is denoted by $(-C)$.

For any PCA (B, C), we define a binary relation " $\ll_C$" on B (called *non-tangential inclusion*) by

(1) $\quad a \ll_C b \leftrightarrow a(-C)b^*$.

Sometimes we will write simply " $\ll$" instead of " $\ll_C$".

We will also consider precontact algebras satisfying some additional axioms:

$(Cref)$ If $a \neq 0$ then aCa (reflexivity axiom);

$(Csym)$ If aCb then bCa (symmetry axiom);

(Ctr) If $a \ll_C c$ then $(\exists b)(a \ll_C b \ll_C c)$ (transitivity axiom);

$(Ccon)$ If $a \neq 0, 1$ then aCa^* or a^*Ca (connectedness axiom).

A precontact algebra (B, C) is called a *contact algebra* ([13]) (and C is called a *contact relation*) if it satisfies the axioms $(Cref)$ and $(Csym)$. We say that two contact algebras are *CA-isomorphic* if they are PCA-isomorphic; also, a PCA-isomorphism between two contact algebras will be called a *CA-isomorphism*.

A precontact algebra (B, C) is called *connected* if it satisfies the axiom $(Ccon)$.

Lemma 2.2. *Let (B, C) be a precontact algebra. Define*

$$aC^{\#}b \iff ((aCb) \vee (bCa) \vee (a.b \neq 0)).$$

Then $C^{\#}$ is a contact relation on B and hence $(B, C^{\#})$ is a contact algebra.

Remark 2.3. We will also consider precontact algebras satisfying the following variant of the transitivity axiom (Ctr):

$(Ctr\#)$ If $a \ll_{C^{\#}} c$ then $(\exists b)(a \ll_{C^{\#}} b \ll_{C^{\#}} c)$.

The axiom $(Ctr\#)$ is known as the "Interpolation axiom".

A contact algebra (B, C) is called a *normal contact algebra* ([6, 21]) if it satisfies the axiom $(Ctr\#)$ and the following one:

(C6) If $a \neq 1$ then there exists $b \neq 0$ such that $b(-C)a$.

The notion of a normal contact algebra was introduced by Fedorchuk [21] (under the name of "*Boolean δ-algebra*") as an equivalent expression of the notion of a *compingent Boolean algebra* of de Vries [6] (see its definition below). We call such algebras "normal contact algebras" because they form a subclass of the class of contact algebras and naturally arise in normal Hausdorff spaces.

The relations C and $\ll$ are inter-definable. For example, normal contact algebras could be equivalently defined (and exactly in this way they were introduced (under the name of *compingent Boolean algebras*) by de Vries in [6]) as a pair of a

Boolean algebra $B = (B, 0, 1, +, ., ^*)$ and a binary relation $\ll$ on B subject to the following axioms:

($\ll$1) $a \ll b$ implies $a \leq b$;
($\ll$2) $0 \ll 0$;
($\ll$3) $a \leq b \ll c \leq t$ implies $a \ll t$;
($\ll$4) ($a \ll b$ and $a \ll c$) implies $a \ll b.c$;
($\ll$5) If $a \ll c$ then $a \ll b \ll c$ for some $b \in B$;
($\ll$6) If $a \neq 0$ then there exists $b \neq 0$ such that $b \ll a$;
($\ll$7) $a \ll b$ implies $b^* \ll a^*$.

Note that if $0 \neq 1$ then the axiom ($\ll$2) follows from the axioms ($\ll$3), ($\ll$4), ($\ll$6) and ($\ll$7).

Obviously, contact algebras could be equivalently defined as a pair of a Boolean algebra B and a binary relation $\ll$ on B subject to the axioms ($\ll$1)-($\ll$4) and ($\ll$7); then, clearly, the relation $\ll$ satisfies also the axioms

($\ll$2') $1 \ll 1$;
($\ll$4') ($a \ll c$ and $b \ll c$) implies $(a + b) \ll c$.

It is not difficult to see that precontact algebras could be equivalently defined as a pair of a Boolean algebra B and a binary relation $\ll$ on B subject to the axioms ($\ll$2), ($\ll$2'), ($\ll$3), ($\ll$4) and ($\ll$4').

It is easy to see that axiom (C6) can be stated equivalently in the form of ($\ll$6).

Now we will give some examples of precontact and contact algebras. We will start with the *extremal contact relations*.

Example 2.4. Let B be a Boolean algebra. Then there exist a largest and a smallest contact relations on B; the largest one, ρ_l (sometimes we will write ρ_l^B), is defined by

$$a \rho_l b \iff (a \neq 0 \text{ and } b \neq 0),$$

and the smallest one, ρ_s (sometimes we will write ρ_s^B), by

$$a \rho_s b \iff a.b \neq 0.$$

Note that, for $a, b \in B$,

$$a \ll_{\rho_s} b \iff a \leq b;$$

hence $a \ll_{\rho_s} a$, for any $a \in B$. Thus (B, ρ_s) is a normal contact algebra.

We are now going to recall the definition of an adjacency space ([22], [17]) and the fact that each adjacency space generates canonically a precontact algebra ([17]).

By an *adjacency space* (see [22] and [17]) we mean a relational system (W, R), where W is a non-empty set whose elements are called *cells*, and R is a binary relation on W called the *adjacency relation*; the subsets of W are called *regions*.

The reflexive and symmetric closure $R^\flat$ of R is defined as follows:

(2) $\quad xR^\flat y \iff ((xRy) \vee (yRx) \vee (x = y))$.

A *precontact relation* C_R between the regions of an adjacency space (W, R) is defined as follows: for every $M, N \subseteq W$,

(3) MC_RN iff $(\exists x \in M)(\exists y \in N)(xRy)$.

Proposition 2.5. ([17]) *Let (W, R) be an adjacency space and let 2^W be the Boolean algebra of all subsets of W. Then:*

(a) *$(2^W, C_R)$ is a precontact algebra;*

(b) *$(2^W, C_R)$ is a contact algebra iff R is a reflexive and symmetric relation on W. If R is a reflexive and symmetric relation on W then C_R coincides with $(C_R)^\#$ and $C_{R^\flat}$;*

(c) *C_R satisfies the axiom (Ctr) iff R is a transitive relation on W;*

(d) *C_R satisfies the axiom $(Ccon)$ iff R is a connected relation on W (which means that if $x, y \in W$ and $x \neq y$ then there is an R-path from x to y or from y to x).*

Clearly, Proposition 2.5(a) implies that if B is a Boolean subalgebra of the Boolean algebra 2^W, then (B, C_R) is also a precontact algebra (here (and further on), for simplicity, we denote again by C_R the restriction of the relation C_R to B).

We recall as well that every topological space generates canonically a contact algebra.

2.6. Let X be a topological space and let $RC(X)$ be the set of all regular closed subsets of X (recall that a subset F of X is said to be *regular closed* if $F = \mathrm{cl}(\mathrm{int}(F))$). Let us equip $RC(X)$ with the following Boolean operations and *contact relation* C_X:

- $F + G = F \cup G$;
- $F^* = \mathrm{cl}(X \setminus F)$;
- $F.G = \mathrm{cl}(\mathrm{int}(F \cap G))(= (F^* \cup G^*)^*)$;
- $0 = \emptyset$, $1 = X$;
- FC_XG iff $F \cap G \neq \emptyset$.

The following lemma is a well-known fact.

Lemma 2.7. *Let X be a topological space. Then*

$$(RC(X), C_X) = (RC(X), 0, 1, +, ., *, C_X)$$

is a complete contact algebra.

The contact algebras of the type $(RC(X), C_X)$, where X is a topological space, are called *standard contact algebras*.

Definition 2.8. ([15, 16]) Let $\underline{B} = (B, C)$ be a precontact algebra and let U_1, U_2 be ultrafilters of B. We set

(4) $U_1 R_{\underline{B}} U_2$ iff $(\forall a \in U_1)(\forall b \in U_2)(aCb)$ (i.e., iff $U_1 \times U_2 \subseteq C$).

The relational system $(Ult(B), R_{\underline{B}})$ is called the *canonical adjacency space of* $\underline{B}$.

We say that U_1, U_2 are *connected* iff $U_1 (R_{\underline{B}})^\flat U_2$ (see (2) for the notation $R^\flat$).

Definition 2.9. ([15, 16]) Let X be a non-empty topological space and R be a binary relation on X. Then the pair $(CO(X), C_R)$ (see (3) for C_R) is a precontact algebra (by Proposition 2.5(a)), called the *canonical precontact algebra of the relational system* (X, R).

Definition 2.10. ([15, 16]) An adjacency space (X, R) is called a *topological adjacency space* (abbreviated as TAS) if X is a topological space and R is a closed relation on X. When X is a compact Hausdorff zero-dimensional space (i.e., when X is a *Stone space*), we say that the topological adjacency space (X, R) is a *Stone adjacency space*.

Two topological adjacency spaces (X, R) and (X_1, R_1) are said to be *TAS-isomorphic* if there exists a homeomorphism $f : X \longrightarrow X_1$ such that, for every $x, y \in X$, xRy iff $f(x) R_1 f(y)$.

Recall that:
(a) a topological space X is called a *Stone space* if it is a compact Hausdorff zero-dimensional space;
(b) *the Stone space* $S(A)$ *of a Boolean algebra* A is the set $X = Ult(A)$ endowed with a topology $\mathcal{T}$ having as a closed base the family $\{s_A(a) \mid a \in A\}$, where

(5) $s_A(a) = \{u \in X \mid a \in u\}$,

for every $a \in A$; then

$$S(A) = (X, \mathcal{T})$$

is a compact Hausdorff zero-dimensional space (i.e., it is indeed a Stone space), $s_A(A) = CO(X)$ and *the Stone map*

(6) $s_A : A \longrightarrow CO(X), \quad a \mapsto s_A(a)$,

is a Boolean isomorphism; also, the family $\{s_A(a) \mid a \in A\}$ is an open base of $(X, \mathcal{T})$. Further, for every Stone space X and for every $x \in X$, we set

(7) $u_x = \{P \in CO(X) \mid x \in P\}$

(sometimes we will also write u_x^X instead of u_x). Then $u_x \in Ult(CO(X))$ and the map

$$f : X \longrightarrow S(CO(X)), \quad x \mapsto u_x,$$

is a homeomorphism.

When $\underline{B} = (B, C)$ is a precontact algebra, the pair $(S(B), R_{\underline{B}})$ is said to be *the canonical Stone adjacency space of* $\underline{B}$.

Theorem 2.11. ([15, 16]) (a) *Each precontact algebra* $\underline{B} = (B, C)$ *is isomorphic to the canonical precontact algebra* $(CO(X, \mathfrak{T}), C_{R_{\underline{B}}})$ *of the Stone adjacency space* $((X, \mathfrak{T}), R_{\underline{B}})$, *where* $(X, \mathfrak{T}) = S(B)$ *and for every* $u, v \in X$, $uR_{\underline{B}}v \iff u \times v \subseteq C$; *the isomorphism between them is just the Stone map* $s_B : B \longrightarrow CO(X, \mathfrak{T})$. *Moreover, the relation* C *satisfies the axiom (Cref) (resp., (Csym); (Ctr)) iff the relation* $R_{\underline{B}}$ *is reflexive (resp., symmetric; transitive).*

(b) *There exists a bijective correspondence between the class of all, up to PCA-isomorphism, precontact algebras and the class of all, up to TAS-isomorphism, Stone adjacency spaces* (X, R); *namely, for each precontact algebra* $\underline{B} = (B, C)$, *the PCA-isomorphism class* $[\underline{B}]$ *of* $\underline{B}$ *corresponds to the TAS-isomorphism class of the canonical Stone adjacency space* $(S(B), R_{\underline{B}})$ *of* $\underline{B}$, *and for each Stone adjacency space* (X, R), *the TAS-isomorphism class* $[(X, R)]$ *of* (X, R) *corresponds to the PCA-isomorphism class of the canonical precontact algebra* $(CO(X), C_R)$ *of* (X, R) *(see (3) for* C_R*).*

Let us also recall the following well known statement (see, e.g., [5], p.271).

Lemma 2.12. *Let* X *be a dense subspace of a topological space* Y. *Then the functions*

$$r : RC(Y) \longrightarrow RC(X),\ F \mapsto F \cap X,$$

and

$$e : RC(X) \longrightarrow RC(Y),\ G \mapsto \mathrm{cl}_Y(G),$$

are Boolean isomorphisms between Boolean algebras $RC(X)$ *and* $RC(Y)$, *and* $e \circ r = id_{RC(Y)}$, $r \circ e = id_{RC(X)}$. *(We will sometimes write* $r_{X,Y}$ *(resp.,* $e_{X,Y}$*) instead of* r *(resp.,* e*).)*

Definition 2.13. ([15, 16]) (a) Let X be a topological space and X_0 be a dense subspace of X. Then the pair (X, X_0) is called a *topological pair*.

(b) Let (X, X_0) be a topological pair. Then we set

(8) $\quad RC(X, X_0) = \{cl_X(A) \mid A \in CO(X_0)\}.$

Lemma 2.14. ([15, 16]) *Let* (X, X_0) *be a topological pair. Then* $RC(X, X_0) \subseteq RC(X)$; *the set* $RC(X, X_0)$ *with the standard Boolean operations on the regular closed subsets of* X *is a Boolean subalgebra of* $RC(X)$; $RC(X, X_0)$ *is isomorphic to the Boolean algebra* $CO(X_0)$; *the sets* $RC(X)$ *and* $RC(X, X_0)$ *coincide iff* X_0 *is an extremally disconnected space. If*

$$C_{(X,X_0)}$$

is the restriction of the contact relation C_X *(see Lemma 2.7) to* $RC(X, X_0)$, *then* $(RC(X, X_0), C_{(X,X_0)})$ *is a contact subalgebra of* $(RC(X), C_X)$.

Definition 2.15. Let $\underline{B} = (B, C)$ be a precontact algebra. A non-empty subset Γ of B is called a *clan* if it satisfies the following conditions:

$(Clan1) \quad 0 \notin \Gamma;$

($Clan2$) If $a \in \Gamma$ and $a \leq b$ then $b \in \Gamma$;

($Clan3$) If $a + b \in \Gamma$ then $a \in \Gamma$ or $b \in \Gamma$;

($Clan4$) If $a, b \in \Gamma$ then $aC^{\#}b$.

The set of all clans of a precontact algebra $\underline{B}$ is denoted by $Clans(\underline{B})$.

Recall that a non-empty subset of a Boolean algebra B is called a *grill* if it satisfies the axioms (Clan1)-(Clan3). The set of all grills of B will be denoted by $Grills(B)$.

We will need the following well-known lemma (see, e.g., [28]):

Lemma 2.16. (Grill Lemma.) *If F is a filter of a Boolean algebra B and G is a grill of B such that $F \subseteq G$ then there exists an ultrafilter U of B with $F \subseteq U \subseteq G$.*

Notation 2.17. Let $(X, \mathcal{T})$ be a topological space, X_0 be a subspace of X, $x \in X$ and B be a subalgebra of the Boolean algebra $(RC(X), +, ., *, \emptyset, X)$ defined in 2.6. We put

(9) $\sigma_x^B = \{F \in B \mid x \in F\}$; $\Gamma_{x,X_0} = \{F \in CO(X_0) \mid x \in cl_X(F)\}$.

We set also

(10) $\nu_x^B = \{F \in B \mid x \in \mathrm{int}_X(F)\}$.

When $B = RC(X)$, we will often write simply σ_x and ν_x instead of, respectively, σ_x^B and ν_x^B; in this case we will sometimes use the notation σ_x^X and ν_x^X as well.

Definition 2.18. (2-Precontact spaces.)([15, 16])
(a) A triple $\underline{X} = (X, X_0, R)$ is called a *2-precontact space* (abbreviated as PCS) if the following conditions are satisfied:

($PCS1$) (X, X_0) is a topological pair and X is a T_0-space;

($PCS2$) (X_0, R) is a Stone adjacency space;

($PCS3$) $RC(X, X_0)$ is a closed base for X;

($PCS4$) For every $F, G \in CO(X_0)$, $\mathrm{cl}_X(F) \cap \mathrm{cl}_X(G) \neq \emptyset$ implies that $F(C_R)^{\#}G$ (see (3) for C_R);

($PCS5$) If $\Gamma \in Clans(CO(X_0), C_R)$ then there exists a point $x \in X$ such that $\Gamma = \Gamma_{x,X_0}$ (see (9) for Γ_{x,X_0}).

(b) Let $\underline{X} = (X, X_0, R)$ be a 2-precontact space. Define, for every $F, G \in RC(X, X_0)$,

$$F \; C_{\underline{X}} \; G \iff ((\exists x \in F \cap X_0)(\exists y \in G \cap X_0)(xRy)).$$

Then the precontact algebra

$$\underline{B}(\underline{X}) = (RC(X, X_0), C_{\underline{X}})$$

is said to be the *canonical precontact algebra of* $\underline{X}$.

(c) A 2-precontact space $\underline{X} = (X, X_0, R)$ is called *reflexive* (resp., *symmetric*; *transitive*) if the relation R is reflexive (resp., symmetric; transitive); $\underline{X}$ is called *connected* if the space X is connected.

(d) Let $\underline{X} = (X, X_0, R)$ and $\underline{\widehat{X}} = (\widehat{X}, \widehat{X}_0, \widehat{R})$ be two 2-precontact spaces. We say that $\underline{X}$ and $\underline{\widehat{X}}$ are *PCS-isomorphic* (or, simply, *isomorphic*) if there exists a homeomorphism $f : X \longrightarrow \widehat{X}$ such that:

(ISO1) $\quad f(X_0) = \widehat{X}_0$; and

(ISO2) $\quad (\forall x, y \in X_0)(xRy \leftrightarrow f(x)\widehat{R}f(y))$.

Remark 2.19. ([15, 16]) It is very easy to see that the canonical precontact algebra of a 2-precontact space, defined in Definition 2.18(b), is indeed a precontact algebra.

Definition 2.20. ([15, 16]) Let $\underline{B} = (B, C)$ be a precontact algebra. We associate with $\underline{B}$ a 2-precontact space

$$\underline{X}(\underline{B}) = (X, X_0, R),$$

called the *canonical 2-precontact space of* $\underline{B}$, as follows:

- $X = Clans(\underline{B})$ and $X_0 = Ult(B)$;
- The topology $\mathfrak{T}$ on the set X is defined in the following way: the family

  $$\{g_{\underline{B}}(a) \mid a \in B\},$$

  where, for any $a \in B$,

  (11) $\quad g_{\underline{B}}(a) = \{\Gamma \in X \mid a \in \Gamma\}$,

  is a closed base of $\mathfrak{T}$. The topology on X_0 is the subspace topology induced by $(X, \mathfrak{T})$.
- $R = R_{\underline{B}}$ (see (4) for the notation $R_{\underline{B}}$).

Proposition 2.21. ([15, 16]) *Let $\underline{B} = (B, C)$ be a precontact algebra. Then the canonical 2-precontact space $\underline{X}(\underline{B}) = (X, X_0, R)$ of $\underline{B}$ defined above is indeed a 2-precontact space.*

Theorem 2.22. (Representation theorem for precontact algebras.)([15, 16])

(a) *Let $\underline{B} = (B, C)$ be a precontact algebra and let $\underline{X}(\underline{B}) = (X, X_0, R)$ be the canonical 2-precontact space of $\underline{B}$. Then the function $g_{\underline{B}} : (B, C) \longrightarrow 2^X$, defined in (11), is a PCA-isomorphism from (B, C) onto the canonical precontact algebra $(RC(X, X_0), C_{\underline{X}(\underline{B})})$ of $\underline{X}(\underline{B})$. The same function $g_{\underline{B}}$ is a PCA-isomorphism between contact algebras $(B, C^{\#})$ and $(RC(X, X_0), C_{(X,X_0)})$ (see Lemma 2.14(a) for $C_{(X,X_0)}$). The sets $RC(X)$ and $RC(X, X_0)$ coincide iff the precontact algebra $\underline{B}$ is complete. The algebra $\underline{B}$ satisfies the axiom $(Cref)$ (resp., $(Csym)$; (Ctr)) iff the 2-precontact space $\underline{X}(\underline{B})$ is reflexive (resp., symmetric; transitive). The algebra $\underline{B}$ is connected iff $\underline{X}(\underline{B})$ is connected.*

(b) *There exists a bijective correspondence between the class of all, up to PCA-isomorphism, (connected) precontact algebras and the class of all, up to PCS-isomorphism, (connected) 2-precontact spaces; namely, for every precontact algebra* $\underline{B}$*, the PCA-isomorphism class* $[\underline{B}]$ *of* $\underline{B}$ *corresponds to the PCS-isomorphism class* $[\underline{X}(\underline{B})]$ *of the canonical 2-precontact space* $\underline{X}(\underline{B})$ *of* $\underline{B}$*, and for every 2-precontact space* $\underline{X}$*, the PCS-isomorphism class* $[\underline{X}]$ *of* $\underline{X}$ *corresponds to the PCA-isomorphism class* $[\underline{B}(\underline{X})]$ *of the canonical precontact algebra* $\underline{B}(\underline{X})$ *of* $\underline{X}$*.*

Corollary 2.23. ([16]) *If* $\underline{X} = (X, X_0, R)$ *is a 2-precontact space then* X *is a compact space.*

Proposition 2.24. ([15, 16]) *Let* X_0 *be a subspace of a topological space* X*. For every* $F, G \in CO(X_0)$*, set*

(12) $F\delta_{(X,X_0)}G$ *iff* $cl_X(F) \cap cl_X(G) \neq \emptyset$.

Then $(CO(X_0), \delta_{(X,X_0)})$ *is a contact algebra.*

Definition 2.25. (2-Contact spaces.)([15, 16]) (a) A topological pair (X, X_0) is called a *2-contact space* (abbreviated as CS) if the following conditions are satisfied:

$(CS1)$ X is a T_0-space;

$(CS2)$ X_0 is a Stone space;

$(CS3)$ $RC(X, X_0)$ is a closed base for X;

$(CS4)$ If $\Gamma \in Clans(CO(X_0), \delta_{(X,X_0)})$ (see (12) for the notation $\delta_{(X,X_0)}$) then there exists a point $x \in X$ such that $\Gamma = \Gamma_{x,X_0}$ (see (9) for Γ_{x,X_0}).

A 2-contact space (X, X_0) is called *connected* if the space X is connected.

(b) Let (X, X_0) be a 2-contact space. Then the contact algebra

$$\underline{B}^c(X, X_0) = (RC(X, X_0), C_{(X,X_0)})$$

(see Lemma 2.14(a) for the notation $C_{(X,X_0)}$) is said to be the *canonical contact algebra of the 2-contact space* (X, X_0).

(c) Let $\underline{B} = (B, C)$ be a contact algebra, $X = Clans(B, C)$, $X_0 = Ult(B)$ and $\mathfrak{T}$ be the topology on X described in Definition 2.20. Take the subspace topology on X_0. Then the pair

$$\underline{X}^c(\underline{B}) = (X, X_0)$$

is called the *canonical 2-contact space of the contact algebra* (B, C). (Note that $\underline{X}^c(\underline{B})$ is indeed a 2-contact space (see [16]).)

(d) Let (X, X_0) and $(\widehat{X}, \widehat{X}_0)$ be two 2-contact spaces. We say that (X, X_0) and $(\widehat{X}, \widehat{X}_0)$ are *CS-isomorphic* (or, simply, *isomorphic*) if there exists a homeomorphism $f : X \longrightarrow \widehat{X}$ such that $f(X_0) = \widehat{X}_0$.

Corollary 2.26. ([16]) *If* $\underline{X} = (X, X_0, R)$ *is a 2-contact space then* X *is a compact space.*

Definition 2.27. (STONE 2-SPACES.)([16])
(a) A topological pair (X, X_0) is called a *Stone 2-space* (abbreviated as S2S) if it satisfies conditions (CS1)-(CS3) of Definition 2.25 and the following condition:

$(S2S4)$ If $\Gamma \in Grills(CO(X_0))$ then there exists a point $x \in X$ such that $\Gamma = \Gamma_{x,X_0}$ (see (9) for Γ_{x,X_0}).

(b) Let (X, X_0) and $(\widehat{X}, \widehat{X}_0)$ be two Stone 2-spaces. We say that (X, X_0) and $(\widehat{X}, \widehat{X}_0)$ are *S2S-isomorphic* (or, simply, *isomorphic*) if there exists a homeomorphism $f : X \longrightarrow \widehat{X}$ such that $f(X_0) = \widehat{X}_0$.

Definition 2.28. ([16]) Let (X, X_0) and (X', X_0') be two Stone 2-spaces and $f : X \longrightarrow X'$ be a continuous map. Then f is called a *2-map* if $f(X_0) \subseteq X_0'$.

The category of all Stone 2-spaces and all 2-maps between them will be denoted by **2Stone**.

The category of all Boolean algebras and all Boolean homomorphisms between them will be denoted by **Bool**.

Recall that a topological space X is said to be *semiregular* if $RC(X)$ is a closed base for X.

Definition 2.29. ([13]) A semiregular T_0-space $(X, \mathfrak{T})$ is said to be *C-semiregular* if for every clan Γ in $(RC(X), C_X)$ there exists a point $x \in X$ such that $\Gamma = \sigma_x$ (see (9) for σ_x).

Proposition 2.30. ([13, Fact 4.1]) *Every C-semiregular space* X *is a compact space.*

Definition 2.31. ([16]) Let $(X, \mathfrak{T})$ be a topological space and $x \in X$. The point x is said to be an *u-point* if for every $U, V \in \mathfrak{T}$, $x \in \mathrm{cl}(U) \cap \mathrm{cl}(V)$ implies that $x \in \mathrm{cl}(U \cap V)$.

Theorem 2.32. ([16]) *If* X *is C-semiregular and*

$$X_0 = \{x \in X \mid x \text{ is an u-point of } X\},$$

then the pair (X, X_0) *is a 2-contact space and* X_0 *is a dense extremally disconnected compact Hausdorff subspace of* X*; moreover,* X_0 *is the unique dense extremally disconnected compact Hausdorff subspace of* X*.*

Lemma 2.33. ([16]) *If* (X, X_0) *is a 2-contact space and* X_0 *is extremally disconnected, then* X *is C-semiregular.*

3 The Main Theorem and its corollaries

The next lemma is obvious.

Lemma 3.1. *Let A be a subalgebra of a Boolean algebra B and (B, C) be a precontact algebra. Then $(A, C \cap A^2)$ is a precontact algebra and if $\Gamma \in Clans(B, C)$ then $\Gamma \cap A \in Clans(A, C \cap A^2)$.*

We will need also a lemma from [16, 13].

Lemma 3.2. ([16, 13]) *Let X be a topological space, B be a subalgebra of the Boolean algebra $(RC(X), +, ., *, \emptyset, X)$ (defined in 2.6), Γ be a grill of B, $x \in X$ and $\Gamma \subseteq \sigma_x^B$. Then $\nu_x^B \subseteq \Gamma$ (see (10) for ν_x^B and (9) for σ_x^B).*

Definition 3.3. Let **PCA** be the category of all precontact algebras and all Boolean homomorphisms $\varphi : (B, C) \longrightarrow (B', C')$ between them such that, for all $a, b \in B$, $\varphi(a)C'\varphi(b)$ implies that aCb.

Let **PCS** be the category of all 2-precontact spaces and all continuous maps $f : (X, X_0, R) \longrightarrow (X', X_0', R')$ between them such that $f(X_0) \subseteq X_0'$ and, for every $x, y \in X_0$, xRy implies that $f(x)R'f(y)$.

The proof of the next theorem is similar in some of its parts to the proof of [16, Theorem 8.17] but, for convenience of the reader, we give here all necessary details.

Theorem 3.4. (The Main Theorem: A Duality Theorem for Precontact Algebras) *The categories* **PCA** *and* **PCS** *are dually equivalent. In fact, the contravariant functor $G^a : \mathbf{PCA} \longrightarrow \mathbf{PCS}$ defined by $G^a(\underline{B}) = \underline{X}(\underline{B})$ on the objects $\underline{B}$ of the category* **PCA**, *and by $G^a(\varphi) : G^a(\underline{B'}) \longrightarrow G^a(\underline{B})$, $\Gamma \mapsto \varphi^{-1}(\Gamma)$, for every* **PCA***-morphism $\varphi : \underline{B} \longrightarrow \underline{B'}$, is a duality functor.*

Proof. We will first define two contravariant functors

$$G^a : \mathbf{PCA} \longrightarrow \mathbf{PCS} \quad \text{and} \quad G^t : \mathbf{PCS} \longrightarrow \mathbf{PCA}.$$

Let $\underline{X} = (X, X_0, R) \in |\mathbf{PCS}|$. Define

$$G^t(X, X_0, R) = \underline{B}(\underline{X}),$$

i.e. $G^t(X, X_0, R)$ is the canonical precontact algebra $\underline{B}(\underline{X}) = (RC(X, X_0), C_{\underline{X}})$ of the 2-precontact space $\underline{X}$ (see Definition 2.18(b) and recall that for every $F_1, F_2 \in RC(X, X_0)$,

$$F_1 C_{\underline{X}} F_2 \iff ((\exists x_1 \in F_1 \cap X_0)(\exists x_2 \in F_2 \cap X_0)(x_1 R x_2))).$$

Hence $G^t(X, X_0, R) \in |\mathbf{PCA}|$.

Let $f \in \mathbf{PCS}((X, X_0, R), (Y, Y_0, R'))$. Define

$$G^t(f) : G^t(Y, Y_0, R') \longrightarrow G^t(X, X_0, R)$$

by the formula

(13) $G^t(f)(\mathrm{cl}_Y(F')) = \mathrm{cl}_X(X_0 \cap f^{-1}(F')), \quad \forall F' \in CO(Y_0).$

Set $\varphi_f = G^t(f)$, $\underline{X} = (X, X_0, R)$, $\underline{Y} = (Y, Y_0, R')$, $C = C_{\underline{X}}$ and $C' = C_{\underline{Y}}$. We will show that φ_f is a Boolean homomorphism between the Boolean algebras $RC(Y, Y_0)$ and $RC(X, X_0)$. Clearly, for every $F' \in CO(Y_0)$, we have that $X_0 \cap f^{-1}(F') = (f|_{X_0})^{-1}(F') \in CO(X_0)$. Hence $\varphi_f(RC(Y, Y_0)) \subseteq RC(X, X_0)$. Obviously, $\varphi_f(\emptyset) = \emptyset$ and $\varphi_f(Y) = X$. Let $F', G' \in CO(Y_0)$. Then $\varphi_f(\mathrm{cl}_Y(F') + \mathrm{cl}_Y(G')) = \varphi_f(\mathrm{cl}_Y(F' \cup G')) = \mathrm{cl}_X(X_0 \cap f^{-1}(F' \cup G')) = \mathrm{cl}_X((X_0 \cap f^{-1}(F')) \cup (X_0 \cap f^{-1}(G'))) = \varphi_f(\mathrm{cl}_Y(F')) + \varphi_f(\mathrm{cl}_Y(G'))$. Also, using Lemma 2.12, we get that $\varphi_f((\mathrm{cl}_Y(F'))^*) = \varphi_f(\mathrm{cl}_Y(Y_0 \setminus F') = \mathrm{cl}_X(X_0 \cap f^{-1}(Y_0 \setminus F')) = \mathrm{cl}_X(X_0 \cap (f^{-1}(Y_0) \setminus f^{-1}(F'))) = \mathrm{cl}_X(X_0 \setminus (X_0 \cap f^{-1}(F'))) = (\mathrm{cl}_X(X_0 \cap f^{-1}(F')))^* = (\varphi_f(\mathrm{cl}_Y(F'))^*$. So,

(14) $G^t(f)$ is a Boolean homomorphism.

Let $F', G' \in CO(Y_0)$ and $\varphi_f(\mathrm{cl}_Y(F'))C\varphi_f(\mathrm{cl}_Y(G'))$. Then

$$\mathrm{cl}_X(X_0 \cap f^{-1}(F'))C\mathrm{cl}_X(X_0 \cap f^{-1}(G'))$$

and thus there exist $x \in X_0 \cap \mathrm{cl}_X(X_0 \cap f^{-1}(F'))(= X_0 \cap f^{-1}(F'))$ and $y \in X_0 \cap \mathrm{cl}_X(X_0 \cap f^{-1}(G'))(= X_0 \cap f^{-1}(G'))$ such that xRy. Hence $f(x)R'f(y)$. By Lemma 2.12, we get that $F' = Y_0 \cap \mathrm{cl}_Y(F')$ and $G' = Y_0 \cap \mathrm{cl}_Y(G')$. Since, obviously, $f(x) \in F'$ and $f(y) \in G'$, we get that $\mathrm{cl}_Y(F')C'\mathrm{cl}_Y(G')$. Thus φ is a **PCA**-morphism, i.e., $G^t(f)$ is well defined.

Now we will show that G^t is a contravariant functor. Clearly, $G^t(id_{(X,X_0,R)}) = id_{G^t(X,X_0,R)}$.

Let $f \in \mathbf{PCS}((X, X_0, R), (Y, Y_0, R'))$ and $g \in \mathbf{PCS}((Y, Y_0, R'), (Z, Z_0, R''))$. Then, for every $F \in CO(Z_0)$, $G^t(g \circ f)(\mathrm{cl}_Z(F)) = \mathrm{cl}_X(X_0 \cap (g \circ f)^{-1}(F)) = \mathrm{cl}_X(X_0 \cap f^{-1}(g^{-1}(F)))$ and $(G^t(f) \circ G^t(g))(\mathrm{cl}_Z(F)) = G^t(f)(\mathrm{cl}_Y(Y_0 \cap g^{-1}(F))) = \mathrm{cl}_X(X_0 \cap f^{-1}(Y_0 \cap g^{-1}(F))) = \mathrm{cl}_X(X_0 \cap f^{-1}(Y_0) \cap f^{-1}(g^{-1}(F))) = \mathrm{cl}_X(X_0 \cap f^{-1}(g^{-1}(F))) = G^t(g \circ f)(\mathrm{cl}_Z(F))$. So, G^t is a contravariant functor.

For every precontact algebra $\underline{B}$, set

$$G^a(\underline{B}) = \underline{X}(\underline{B}),$$

where $\underline{X}(\underline{B})$ is the canonical 2-precontact space of the precontact algebra $\underline{B}$ (see Definition 2.20). Then Proposition 2.21 implies that $G^a(\underline{B}) \in |\mathbf{PCS}|$.

Let $\underline{B} = (B, C)$, $\underline{B}' = (B', C')$ and $\varphi \in \mathbf{PCA}(\underline{B}, \underline{B}')$. Let $G^a(\underline{B}) = (X, X_0, R)$ and $G^a(\underline{B}') = (Y, Y_0, R')$. Then we define the map

$$G^a(\varphi) : G^a(\underline{B}') \longrightarrow G^a(\underline{B})$$

by the formula

(15) $G^a(\varphi)(\Gamma) = \varphi^{-1}(\Gamma), \quad \forall \Gamma \in Y.$

Set $f_\varphi = G^a(\varphi)$. Since every grill of a Boolean algebra B' is a union of ultrafilters of B', every union of ultrafilters of B' is a grill of B' (see, e.g., [13, Corollary 3.1]), and

the inverse image of an ultrafilter by a Boolean homomorphism between two Boolean algebras is again an ultrafilter, we get that $\forall \Gamma \in Y$, $f_\varphi(\Gamma)$ is a grill of B. We have to show that $f_\varphi(\Gamma)$ is a clan. Let $a, b \in f_\varphi(\Gamma)$, i.e. $a, b \in \varphi^{-1}(\Gamma)$. Hence $\varphi(a) \in \Gamma \ni \varphi(b)$. Thus $\varphi(a)(C')^{\#}\varphi(b)$, i.e. $(\varphi(a)C'\varphi(b)) \vee (\varphi(b)C'\varphi(a)) \vee (\varphi(a).\varphi(b) \neq 0)$. Then $(aCb) \vee (bCa) \vee (a.b \neq 0)$. Therefore, $aC^{\#}b$. This shows that $f_\varphi(\Gamma)$ is a clan, i.e. $f_\varphi : Y \longrightarrow X$.

We will show that f_φ is a continuous function. Let $a \in B$. Then $g_{\underline{B}}(a) = \{\Gamma \in X \mid a \in \Gamma\}$ is a basic closed subset of X (see Definition 2.20). We will show that

(16) $f_\varphi^{-1}(g_{\underline{B}}(a)) = g_{\underline{B'}}(\varphi(a))(= \{\Gamma' \in Y \mid \varphi(a) \in \Gamma'\})$.

Indeed, let $\Gamma' \in f_\varphi^{-1}(g_{\underline{B}}(a))$. Then $f_\varphi(\Gamma') \in g_{\underline{B}}(a)$. Thus $a \in \varphi^{-1}(\Gamma')$, i.e. $\varphi(a) \in \Gamma'$. So, $\Gamma' \in g_{\underline{B'}}(\varphi(a))$. Hence $f_\varphi^{-1}(g_{\underline{B}}(a)) \subseteq g_{\underline{B'}}(\varphi(a))$. Conversely, let $\Gamma' \in g_{\underline{B'}}(\varphi(a))$, i.e. $\varphi(a) \in \Gamma'$. Then $a \in \varphi^{-1}(\Gamma') = f_\varphi(\Gamma')$. Hence $f_\varphi(\Gamma') \in g_{\underline{B}}(a)$. Then $\Gamma' \in f_\varphi^{-1}(g_{\underline{B}}(a))$. So, $f_\varphi^{-1}(g_{\underline{B}}(a)) \supseteq g_{\underline{B'}}(\varphi(a))$. Thus the equation (16) is verified and we get that f_φ is a continuous function.

Let us now show that $f_\varphi(Y_0) \subseteq X_0$. Let $u' \in Y_0$. Then $u' \in Ult(B)$. Hence $f_\varphi(u') = \varphi^{-1}(u') \in Ult(A) = X_0$. Therefore, $f_\varphi(Y_0) \subseteq X_0$.

Let $x, y \in Y_0$ and $xR'y$. Then $x, y \in Ult(B')$. Hence $f_\varphi(x), f_\varphi(y) \in Ult(B)$. We will show that $f_\varphi(x)Rf_\varphi(y)$. We have that $f_\varphi(x)Rf_\varphi(y) \iff f_\varphi(x) \times f_\varphi(y) \subseteq C \iff (\forall a \in \varphi^{-1}(x))(\forall b \in \varphi^{-1}(y))(aCb)$. Now, since $xR'y$, we have that $x \times y \subseteq C'$, i.e. $(\forall a \in x)(\forall b \in y)(aC'b)$. Thus, if $a \in \varphi^{-1}(x)$ and $b \in \varphi^{-1}(y)$, then $\varphi(a) \in x$ and $\varphi(b) \in y$; hence $\varphi(a)C'\varphi(b)$ and, therefore, aCb. This shows that $f_\varphi(x)Rf_\varphi(y)$. So,

$$G^a(\varphi) \in \mathbf{PCS}(G^a(\underline{B'}), G^a(\underline{B})).$$

Clearly, for every precontact algebra (B, C), $G^a(id_{(B,C)}) = id_{G^a(B,C)}$. Let $\varphi \in \mathbf{PCA}((B, C), (B_1, C_1))$ and $\psi \in \mathbf{PCA}((B_1, C_1), (B_2, C_2))$. Let $f_\varphi = G^a(\varphi)$, $f_\psi = G^a(\psi)$ and $G^a(B_2, C_2) = (Z, Z_0, R)$. Then, for every $\Gamma \in Z$, we have that $G^a(\psi \circ \varphi)(\Gamma) = (\psi \circ \varphi)^{-1}(\Gamma) = \varphi^{-1}(\psi^{-1}(\Gamma)) = f_\varphi(f_\psi(\Gamma)) = (G^a(\varphi) \circ G^a(\psi))(\Gamma)$. We get that G^a is a contravariant functor.

Let $\underline{X} = (X, X_0, R) \in |\mathbf{PCS}|$. We will fix a **PCS**-isomorphism $t_{\underline{X}}$ between $\underline{X}$ and $G^a(G^t(\underline{X}))$. We have that $G^t(X, X_0, R) = \underline{B}(\underline{X}) = (RC(X, X_0), C_{\underline{X}})$. Set

$$\underline{B} = (B, C) = (RC(X, X_0), C_{\underline{X}}).$$

Let $G^a(\underline{B}) = (Y, Y_0, R')$. Then $Y = Clans(\underline{B})$, $Y_0 = Ult(B)$ and, for every $x, y \in Y_0$, $xR'y \iff x \times y \subseteq C$. By [16, Proposition 4.7(a)] and Lemma 2.12, the contact algebras

(17) $(CO(X_0), (C_R)^{\#})$ and $(RC(X, X_0), C_{(X,X_0)})$ are isomorphic.

Hence there exists a bijection between the sets

$$Clans(CO(X_0), (C_R)^{\#}) \text{ and } Clans(RC(X, X_0), C_{(X,X_0)}).$$

We have that for every $F, G \in CO(X_0)$,

$$F(C_R)^{\#}G \iff (FC_RG) \vee (GC_RF) \vee (F.G \neq 0) \iff$$

$$\iff [(\exists x \in F)(\exists y \in G)(xRy)] \vee [(\exists y \in G)(\exists x \in F)(yRx)] \vee (F.G \neq 0) \iff$$
$$\iff \mathrm{cl}_X(F)(C_{\underline{X}})^{\#}\mathrm{cl}_X(G).$$

Hence the contact algebras

(18) $(RC(X, X_0), (C_{\underline{X}})^{\#})$, $(CO(X_0), (C_R)^{\#})$ and $(RC(X, X_0), C_{(X,X_0)})$

are isomorphic. In fact, the contact algebras

(19) $(RC(X, X_0), (C_{\underline{X}})^{\#})$ and $(RC(X, X_0), C_{(X,X_0)})$ coincide.

Therefore, the sets

$$Clans(RC(X, X_0), C_{\underline{X}})(= Y) \text{ and } Clans(RC(X, X_0), C_{(X,X_0)})$$

coincide. By [13, Proposition 4.1(ii)], for every $x \in X$, $\sigma_x \in Clans(RC(X), C_X)$. Hence, by Lemma 3.1, for every subalgebra A of the Boolean algebra $RC(X)$,

(20) $\sigma_x^A \in Clans(A, C_X \cap A^2)$.

Therefore, for every $x \in X$,

(21) $\sigma_x^{RC(X,X_0)} \in Clans(RC(X, X_0), C_{(X,X_0)})$.

So, the following map is well-defined:

(22) $t_{\underline{X}} : (X, X_0, R) \longrightarrow G^a(G^t(X, X_0, R))(= (Y, Y_0, R'))$, $x \mapsto \sigma_x^{RC(X,X_0)}$;

we will show that it is a homeomorphism. We start by proving that $t_{\underline{X}}$ is a surjection. Let $\Gamma \in Y$. Then $\Gamma \in Clans(\underline{B})$. From the above considerations, we get that $\Gamma' = r_{X_0,X}(\Gamma) \in Clans(CO(X_0), C_R)$. Hence, by (PCS5), there exists $x \in X$ such that $\Gamma' = \Gamma_{x,X_0}$. Since, by Lemma 2.12, $\Gamma = e_{X_0,X}(\Gamma')$, we get that $\Gamma = \sigma_x^B = t_{\underline{X}}(x)$. So, $t_{\underline{X}}$ is a surjection. For showing that $t_{\underline{X}}$ is a injection, let $x, y \in X$ and $x \neq y$. Since X is a T_0-space, there exists an open subset U of X such that $|U \cap \{x, y\}| = 1$. We can suppose, without loss of generality, that $x \in U$ and $y \notin U$. Since B is a closed base of X, there exists $F \in B$ such that $x \in X \setminus F \subseteq U$. Then $y \in F$ and $x \notin F$. Hence $F \in \sigma_y^B$ and $F \notin \sigma_x^B$, i.e. $t_{\underline{X}}(x) \neq t_{\underline{X}}(y)$. So, $t_{\underline{X}}$ is an injection. Thus $t_{\underline{X}}$ is a bijection. We will now prove that $t_{\underline{X}}$ is a continuous map. We have that the family $\{g_{\underline{B}}(F) = \{\Gamma \in Y \mid F \in \Gamma\} \mid F \in B\}$ is a closed base of Y. Let $F \in B$. We will show that

(23) $t_{\underline{X}}^{-1}(g_{\underline{B}}(F)) = F$.

Let $x \in F$. Set $t_{\underline{X}}(x) = \Gamma$. Then $\Gamma = \sigma_x^B$. Since $F \in \Gamma$, we get that $\Gamma \in g_{\underline{B}}(F)$. Thus $t_{\underline{X}}(F) \subseteq g_{\underline{B}}(F)$, i.e. $F \subseteq t_{\underline{X}}^{-1}(g_{\underline{B}}(F))$. Conversely, let $x \in t_{\underline{X}}^{-1}(g_{\underline{B}}(F))$. Set $\Gamma = t_{\underline{X}}(x)$. Then $\Gamma \in g_{\underline{B}}(F)$. Hence $F \in \Gamma$. Since $\Gamma = \sigma_x^B$, we get that $x \in F$. Hence $F \supseteq t_{\underline{X}}^{-1}(g_{\underline{B}}(F))$. So, $F = t_{\underline{X}}^{-1}(g_{\underline{B}}(F))$. This shows that $t_{\underline{X}}$ is a continuous map. For showing that $t_{\underline{X}}^{-1}$ is a continuous map, let $F \in B$. Using (23) and the fact that $t_{\underline{X}}$ is

a bijection, we get that $t_{\underline{X}}(F) = g_{\underline{B}}(F)$. Hence $(t_{\underline{X}}^{-1})^{-1}(F) = g_{\underline{B}}(F)$. This shows that $t_{\underline{X}}^{-1}$ is a continuous map. So, $t_{\underline{X}}$ is a homeomorphism.

We will now show that $t_{\underline{X}}(X_0) = Y_0$. Let $x \in X_0$. Set $\Gamma = t_{\underline{X}}(x)$. Then $\Gamma = \sigma_x^B$ and $r_{X_0,X}(\Gamma) = \{F \in CO(X_0) \mid x \in F\} = u_x^{X_0} \in Ult(CO(X_0))$. Then, by Lemma 2.12, $\Gamma = e_{X_0,X}(u_x^{X_0}) \in Ult(B) = Y_0$. Hence $t_{\underline{X}}(X_0) \subseteq Y_0$. Let now $\Gamma \in Y_0$. Then $\Gamma \in Ult(B)$ and thus $u = r_{X_0,X}(\Gamma) \in Ult(CO(X_0))$. Clearly, there exist $x \in X_0$ such that $u = u_x^{X_0}$. Then $\Gamma = e_{X_0,X}(u_x^{X_0}) = \sigma_x^B = t_{\underline{X}}(x)$. Therefore, $t_{\underline{X}}(X_0) \supseteq Y_0$. We have proved that $t_{\underline{X}}(X_0) = Y_0$.

Finally, we will prove that for every $x, y \in X_0$, $xRy \iff t_{\underline{X}}(x)R't_{\underline{X}}(y)$. So, let $x, y \in X_0$ and xRy. Then $t_{\underline{X}}(x) = \sigma_x^B$ and $t_{\underline{X}}(y) = \sigma_y^B$. We have that

$$t_{\underline{X}}(x)R't_{\underline{X}}(y) \iff \sigma_x^B \times \sigma_y^B \subseteq C_{\underline{X}} \iff [((x \in F \in B) \wedge (y \in G \in B)) \Rightarrow (FC_{\underline{X}}G)].$$

This shows that, since xRy, $t_{\underline{X}}(x)R't_{\underline{X}}(y)$ holds. Conversely, let $x, y \in X_0$ and $t_{\underline{X}}(x)R't_{\underline{X}}(y)$. Suppose that $x(-R)y$. Then there exist $F, G \in CO(X_0)$ such that $x \in F$, $y \in G$ and $R \cap (F \times G) = \emptyset$. This implies that $\mathrm{cl}_X(F)(-C_{\underline{X}})\mathrm{cl}_X(G)$, a contradiction. Hence, xRy.

All this shows that

(24) $t_{\underline{X}}$ is a **PCS**-isomorphism.

Let $\underline{B} = (B, C)$ be a precontact algebra and let us set $(X, X_0, R) = G^a(\underline{B})$. Then $G^t(X, X_0, R) = (RC(X, X_0), C_{\underline{X}})$ and, using Theorem 2.22(a), we get that the map

$$g_{\underline{B}} : \underline{B} \longrightarrow G^t(G^a(\underline{B}))(= (RC(X, X_0), C_{\underline{X}})), \quad a \mapsto g_{\underline{B}}(a) = \{\Gamma \in X \mid a \in \Gamma\},$$

is a PCA-isomorphism. This implies immediately that $g_{\underline{B}}$ is also a **PCA**-isomorphism.

We will now show that

$$t : Id_{\mathbf{PCS}} \longrightarrow G^a \circ G^t,$$

defined by $t(\underline{X}) = t_{\underline{X}}$, $\forall \underline{X} \in |\mathbf{PCS}|$, is a natural isomorphism.

Let $\underline{X} = (X, X_0, R)$, $\underline{Y} = (Y, Y_0, R')$, $\underline{X}, \underline{Y} \in |\mathbf{PCS}|$, $f \in \mathbf{PCS}(\underline{X}, \underline{Y})$ and $\hat{f} = G^a(G^t(f))$. We will show that

(25) $\hat{f} \circ t_{\underline{X}} = t_{\underline{Y}} \circ f$.

Set $\varphi_f = G^t(f)$, $A = RC(X, X_0)$, $B = RC(Y, Y_0)$, $\underline{B} = G^t(\underline{Y})(= (B, C_{\underline{Y}}))$. Let $x \in X$. Then

$$(t_{\underline{Y}} \circ f)(x) = t_{\underline{Y}}(f(x)) = \sigma_{f(x)}^B = \{F \in B \mid f(x) \in B\}.$$

Further, $\hat{f}(t_{\underline{X}}(x)) = \hat{f}(\sigma_x^A) = \varphi_f^{-1}(\sigma_x^A)$. Set $\Gamma' = \varphi_f^{-1}(\sigma_x^A)$. Then $\Gamma' = \{G \in B \mid \varphi_f(G) \in \sigma_x^A\} = \{\mathrm{cl}_Y(G_0) \mid G_0 \in CO(Y_0), x \in \mathrm{cl}_X(X_0 \cap f^{-1}(G_0))\}$. Let $G_0 \in CO(Y_0)$ and $\mathrm{cl}_Y(G_0) \in \Gamma'$. Then $x \in \mathrm{cl}_X(X_0 \cap f^{-1}(G_0))$ and thus $f(x) \in f(\mathrm{cl}_X(X_0 \cap f^{-1}(G_0)) \subseteq \mathrm{cl}_Y(f(X_0 \cap f^{-1}(G_0))) \subseteq \mathrm{cl}_Y(G_0)$. Therefore,

$$\Gamma' \subseteq \sigma_{f(x)}^B.$$

We have that $\Gamma' \in Clans(\underline{B}) = Clans(RC(Y,Y_0), C_{(Y,Y_0)})$. Hence $\Gamma'_r = r_{Y_0,Y}(\Gamma') \in Clans(CO(Y_0), C_{R'})$. Thus, by (PCS5), there exists $y \in Y$ such that $\Gamma'_r = \Gamma_{y,Y_0}$. Then

(26) $\Gamma' = \sigma^B_y$.

Since $\Gamma' \subseteq \sigma^B_{f(x)}$ and Γ' is a grill, we get, by Lemma 3.2, that $\nu^B_{f(x)} \subseteq \Gamma'$. According to [13, Proposition 4.1], $\nu^B_{f(x)}$ is a filter of B. Hence, by Lemma 2.16, there exists an ultrafilter u of B such that $\nu^B_{f(x)} \subseteq u \subseteq \Gamma'$. Then $u \subseteq \sigma^B_y$ and since u is a grill of B, Lemma 3.2 implies that $\nu^B_y \subseteq u$. So, we obtained that $\nu^B_{f(x)} \cup \nu^B_y \subseteq u \subseteq \Gamma'$. Then, for every $F' \in \nu^B_{f(x)}$ and every $G' \in \nu^B_y$, we have that $F'.G' \neq 0$, i.e. $\mathrm{cl}_Y(\mathrm{int}_Y(F' \cap G')) \neq \emptyset$. Hence $\mathrm{int}_Y(F' \cap G') \neq \emptyset$ and thus $\mathrm{int}_Y(F') \cap \mathrm{int}_Y(G') \neq \emptyset$, for every $F' \in \nu^B_{f(x)}$ and every $G' \in \nu^B_y$. Since Y is a T_0-space, using [16, Lemma 8.14], we get that $y = f(x)$. Therefore $\Gamma' = \sigma^B_{f(x)}$. Thus $\hat{f} \circ t_{\underline{X}} = t_{\underline{Y}} \circ f$ and hence t is a natural isomorphism.

Finally, we will prove that

$$g : Id_{\mathbf{PCA}} \longrightarrow G^t \circ G^a, \text{ where } g(\underline{B}) = g_{\underline{B}}, \quad \forall \underline{B} \in |\mathbf{PCA}|,$$

is a natural isomorphism.

Let $\underline{A} = (A, C)$, $\underline{B} = (B, C')$, $\underline{A}, \underline{B} \in |\mathbf{PCA}|$, $\varphi \in \mathbf{PCA}(\underline{A}, \underline{B})$ and $\hat{\varphi} = G^t(G^a(\varphi))$. We have to prove that $g_{\underline{B}} \circ \varphi = \hat{\varphi} \circ g_{\underline{A}}$. Set $f = G^a(\varphi)$, $(X, X_0, R) = G^a(\underline{A})$ and $(Y, Y_0, R') = G^a(\underline{B})$. Then $\hat{\varphi} = G^t(f) (= \varphi_f)$. Let $a \in A$. Then $g_{\underline{B}}(\varphi(a)) = \{\Gamma' \in Y \mid \varphi(a) \in \Gamma'\}$. Further, using Lemma 2.12, we get that

$$g_{\underline{B}}(\varphi(a)) = \mathrm{cl}_Y(s_B(\varphi(a))) \quad \text{and} \quad g_A(a) = \mathrm{cl}_X(s_A(a))$$

(see (5) for the notation s_A). Thus

$$\hat{\varphi}(g_{\underline{A}}(a)) = \mathrm{cl}_Y(Y_0 \cap f^{-1}(s_A(a))).$$

Let $u' \in Y_0 \cap f^{-1}(s_A(a))$. Then $u' \in Ult(B)$ and $f(u') \in s_A(a)$. Hence $\varphi^{-1}(u') \in s_A(a) = \{u \in Ult(A) \mid a \in u\}$. Thus $a \in \varphi^{-1}(u')$, i.e. $\varphi(a) \in u'$. Therefore $u' \in s_B(\varphi(a))$. So, $Y_0 \cap f^{-1}(s_A(a)) \subseteq s_B(\varphi(a))$. Conversely, let $u' \in s_B(\varphi(a))$. Then $u' \in Y_0$ and $\varphi(a) \in u'$. Hence $a \in \varphi^{-1}(u') = f(u')$. Thus $f(u') \in s_A(a)$. Therefore, $u' \in Y_0 \cap f^{-1}(s_A(a))$. So, $Y_0 \cap f^{-1}(s_A(a)) \supseteq s_B(\varphi(a))$ and we get that $Y_0 \cap f^{-1}(s_A(a)) = s_B(\varphi(a))$. Hence $\hat{\varphi}(g_{\underline{A}}(a)) = \mathrm{cl}_Y(s_B(\varphi(a))) = g_{\underline{B}}(\varphi(a))$. So, g is a natural isomorphism.

We have proved that (G^t, G^a, g, t) is a duality between the categories **PCS** and **PCA**. □

In the proof of the next corollary (as well as in the proofs of all further corollaries of Theorem 3.4), we will use the notation from (the proof of) Theorem 3.4.

Corollary 3.5. *Let $\underline{X} = (X, X_0, R)$, $\underline{Y} = (Y, Y_0, R')$ and $f \in \mathbf{PCS}(\underline{X}, \underline{Y})$. Then, for every $H \in RC(Y, Y_0)$,*

(27) $G^t(f)(H) = f^{-1}(H)$

(see (13) for the notation G^t).

Proof. Recall that $RC(Y, Y_0) = \{\mathrm{cl}_Y(F) \mid F \in CO(Y_0)\}$. Let now $H \in RC(Y, Y_0)$ and $\hat{f} = G^a(G^t(f))$. Then, by (25), $\hat{f} \circ t_{\underline{X}} = t_{\underline{Y}} \circ f$. Thus,

$$f^{-1}(H) = (t_{\underline{X}})^{-1}(\hat{f}^{-1}(t_{\underline{Y}}(H))).$$

Let $\underline{B} = G^t(\underline{X})$, $\underline{B'} = G^t(\underline{Y})$ and $\varphi = G^t(f)$. Using (24), (23) and (16), we get that $t_{\underline{Y}}(H) = g_{\underline{B'}}(H)$ (see (11) for the notation $g_{\underline{B'}}(H)$), $\hat{f}^{-1}(t_{\underline{Y}}(H)) = \hat{f}^{-1}(g_{\underline{B'}}(H)) = g_{\underline{B}}(\varphi(H))$ and $(t_{\underline{X}})^{-1}(\hat{f}^{-1}(t_{\underline{Y}}(H))) = (t_{\underline{X}})^{-1}(g_{\underline{B}}(\varphi(H))) = \varphi(H)$. Hence, $f^{-1}(H) = G^t(f)(H)$. □

As it was already noted, Theorem 3.4 implies the Stone Duality Theorem [27, 24] and its connected version [16].

Corollary 3.6. ([27, 24]) *The category* **Bool** *and the category* **Stone** *of compact zero-dimensional Hausdorff spaces and continuous maps are dually equivalent.*

Proof. Let $\boldsymbol{B}$ be the full subcategory of the category **PCA** having as objects all (pre)contact algebras of the form (B, ρ_s^B) (see Example 2.4 for the notation ρ_s^B). It is easy to see that if A and B are Boolean algebras, then a function $\varphi : A \longrightarrow B$ induces a **PCA**-morphism between (A, ρ_s^A) and (B, ρ_s^B) iff φ is a Boolean homomorphism. Hence the categories **Bool** and $\boldsymbol{B}$ are isomorphic.

Let **S** be the full subcategory of the category **PCS** having as objects all 2-precontact spaces of the form (X, X, D_X), where D_X is the diagonal of X. Obviously, if $(X, X, D_X) \in |\mathbf{S}|$ then X is a Stone space. Conversely, it is easy to see that every triple of the form (X, X, D_X), where X is a Stone space, is a 2-precontact space. Thus the objects of the category **S** are all triples of the form (X, X, D_X), where X is a Stone space. Clearly, if $X, Y \in |\mathbf{Stone}|$, then a function $f : X \longrightarrow Y$ induces a **PCS**-morphism between (X, X, D_X) and (Y, Y, D_Y) iff f is a continuous map. Hence the categories **S** and **Stone** are isomorphic.

We will now show that $G^a(\boldsymbol{B}) \subseteq \mathbf{S}$ and $G^t(\mathbf{S}) \subseteq \boldsymbol{B}$. Indeed, if B is a Boolean algebra, then $G^a(B, \rho_s^B) = (Clans(B, \rho_s^B), Ult(B), R)$, where, for every $u, v \in Ult(B)$, $uRv \iff u \times v \subseteq \rho_s^B$. Obviously, we have that for every $u, v \in Ult(B)$, $uRv \iff u = v$. Further, by [13, Example 3.1], $Clans(B, \rho_s^B) = Ult(B)$. Also, by Theorem 3.4, $G^a(B, \rho_s^B)$ is a 2-precontact space. Hence $G^a(B, \rho_s^B) = (X, X, D_X)$, where $X \in |\mathbf{Stone}|$. So, $G^a(\boldsymbol{B}) \subseteq \mathbf{S}$. Let now $X \in |\mathbf{Stone}|$ and $\underline{X} = (X, X, D_X)$. Then $G^t(\underline{X}) = (RC(X, X), C_{\underline{X}})$. We have that $RC(X, X) = CO(X)$ and for every $F, G \in CO(X)$, $FC_{\underline{X}}G \iff [(\exists x \in F)(\exists y \in G)(xD_Xy)] \iff F \cap G \neq \emptyset \iff F.G \neq 0 \iff F\rho_s^{CO(X)}G$. Thus $G^t(\underline{X}) \in |\mathbf{S}|$. So, we get that $G^t(\mathbf{S}) \subseteq \boldsymbol{B}$. Therefore we obtain, using Theorem 3.4, that the restriction of the contravariant functor G^a to the category $\boldsymbol{B}$ is a duality between the categories $\boldsymbol{B}$ and **S**. This implies that the categories **Bool** and **Stone** are dually equivalent. □

Now we will obtain as a corollary of Theorem 3.4 the connected version of the Stone Duality Theorem proved in [16].

Corollary 3.7. ([16]) *The categories* **Bool** *and* **2Stone** *are dually equivalent.*

Proof. Let $\boldsymbol{B'}$ be the full subcategory of the category **PCA** having as objects all (pre)contact algebras of the form (B, ρ_l^B) (see Example 2.4 for the notation ρ_l^B). It is easy to see that if A and B are Boolean algebras, then a function $\varphi : A \longrightarrow B$ induces a **PCA**-morphism between (A, ρ_l^A) and (B, ρ_l^B) iff φ is a Boolean homomorphism. Hence the categories **Bool** and $\boldsymbol{B'}$ are isomorphic.

Let $\mathbf{S'}$ be the full subcategory of the category **PCS** having as objects all 2-precontact spaces of the form $(X, X_0, (X_0)^2)$. We will show that if $(X, X_0, (X_0)^2) \in |\mathbf{S'}|$ then (X, X_0) is a Stone 2-space. Indeed, set $B = CO(X_0)$ and $R = (X_0)^2$. Then, for every $F, G \in B$, we have that $FC_RG \iff [(\exists x \in F)(\exists y \in G)((x, y) \in R)] \iff [(F \neq \emptyset) \wedge (G \neq \emptyset)] \iff F\rho_l^B G$. Thus, using [13, Example 3.1], we get that $Clans(B, C_R) = Clans(B, \rho_l^B) = Grills(B)$. Now it becomes obvious that the axiom (S2S4) is fulfilled because the axiom (PCS5) is fulfilled. The rest is clear, so that (X, X_0) is a Stone 2-space. Arguing similarly, we get that, conversely, if (X, X_0) is a Stone 2-space then $(X, X_0, (X_0)^2)$ is a 2-precontact space. Hence, the objects of the category $\mathbf{S'}$ are all triples $(X, X_0, (X_0)^2)$, where (X, X_0) is a Stone 2-space (i.e., $(X, X_0) \in |\mathbf{2Stone}|$). Clearly, if $(X, X_0), (Y, Y_0) \in |\mathbf{2Stone}|$, then a function $f : X \longrightarrow Y$ induces a **PCS**-morphism between (X, X_0, X_0^2) and $(Y, Y_0, Y_0^2) \iff (f$ is a continuous map and $f(X_0) \subseteq Y_0) \iff (f \in \mathbf{2Stone}((X, X_0), (Y, Y_0)))$. Hence the categories $\mathbf{S'}$ and **2Stone** are isomorphic.

We will now show that $G^a(\boldsymbol{B'}) \subseteq \mathbf{S'}$ and $G^t(\mathbf{S'}) \subseteq \boldsymbol{B'}$. Indeed, if B is a Boolean algebra, then $G^a(B, \rho_l^B) = (Clans(B, \rho_l^B), Ult(B), R)$, where, for every $u, v \in Ult(B)$, $uRv \iff u \times v \subseteq \rho_l^B$. Since the ultrafilters do not contain the zero element, we get that uRv for every $u, v \in Ult(B)$, i.e. $R = (Ult(B))^2$. Also, by Theorem 3.4, $G^a(B, \rho_l^B)$ is a 2-precontact space. Hence $G^a(B, \rho_l^B) \in |\mathbf{S'}|$. So, $G^a(\boldsymbol{B'}) \subseteq \mathbf{S'}$. Let now $(X, X_0) \in |\mathbf{2Stone}|$ and $\underline{X} = (X, X_0, X_0^2)$. Then $G^t(\underline{X}) = (RC(X, X_0), C_{\underline{X}})$. We have that for every $F, G \in CO(X_0)$, $(\mathrm{cl}_X(F) C_{\underline{X}} \mathrm{cl}_X(G)) \iff [(\exists x \in F)(\exists y \in G)((x, y) \in X_0^2)] \iff [(F \neq \emptyset) \wedge (G \neq \emptyset)] \iff (\mathrm{cl}_X(F) \rho_l^{RC(X,X_0)} \mathrm{cl}_X(G))$. Thus $G^t(\underline{X}) \in |\mathbf{S'}|$. So, we get that $G^t(\mathbf{S'}) \subseteq \boldsymbol{B'}$. Therefore we obtain, using Theorem 3.4, that the restriction of the contravariant functor G^a to the category $\boldsymbol{B'}$ is a duality between the categories $\boldsymbol{B'}$ and $\mathbf{S'}$. This implies that the categories **Bool** and **2Stone** are dually equivalent. □

Recall that: (a)([13]) if (B, C) is a contact algebra, then a non-empty set U of ultrafilters of B is called a *clique in* (B, C) if for any two $u, v \in U$ we have that $u \times v \subseteq C$; (b)([13]) a subset of B is a clan in (B, C) iff it is a union of the elements of a clique in (B, C).

Corollary 3.8. *Let* $\underline{X} = (X, X_0, R_1)$, $\underline{Y} = (Y, Y_0, R_2)$ *and* $\underline{X}, \underline{Y} \in |\mathbf{PCS}|$. *If* $f, g \in \mathbf{PCS}(\underline{X}, \underline{Y})$ *and* $f|_{X_0} = g|_{X_0}$ *then* $f = g$.

Proof. Let $B = RC(X, X_0)$, $B' = RC(Y, Y_0)$ and

$$G^a(G^t(\underline{X})) = (X', X_0', R_1'), \quad G^a(G^t(\underline{Y})) = (Y', Y_0', R_2').$$

Let $f' = G^a(G^t(f))$. Then, by Theorem 3.4, we have that $t_{\underline{Y}} \circ f = f' \circ t_{\underline{X}}$, i.e.

$$f'(\sigma_x^B) = \sigma_{f(x)}^{B'}.$$

Let $x \in X \setminus X_0$. Then $t_{\underline{X}}(x) = \sigma_x^B$ is a clan in $(B, (C_{\underline{X}})^{\#})$ (see (21) and (19)). Hence it is a union of the elements of a clique $\{\sigma_{x_\alpha}^B \mid x_\alpha \in X_0, \alpha \in A\}$ in $(B, (C_{\underline{X}})^{\#})$. Let $\Gamma' = \{\mathrm{cl}_Y(G') \mid G' \in CO(Y_0), f(x_\alpha) \in G' \text{ for some } \alpha \in A\}$. We will show that

(28) $\sigma_{f(x)}^{B'} = \Gamma'$.

Indeed, let $G' \in CO(Y_0)$ and $f(x_\alpha) \in G'$ for some $\alpha \in A$. Set $G = (f|_{X_0})^{-1}(G')$. Then $x_\alpha \in G \in CO(X_0)$. Thus $x \in \mathrm{cl}_X(G)$. Since f is a continuous function, we get that $f(x) \in \mathrm{cl}_Y(f(G)) \subseteq \mathrm{cl}_Y(G')$. Thus $\mathrm{cl}_Y(G') \in \sigma_{f(x)}^{B'}$. Therefore $\Gamma' \subseteq \sigma_{f(x)}^{B'}$. Set, for every $\alpha \in A$, $u_\alpha = t_{\underline{X}}(x_\alpha)(= \sigma_{x_\alpha}^B)$ and $v_\alpha = t_{\underline{Y}}(f(x_\alpha))(= \sigma_{f(x_\alpha)}^{B'})$. Obviously, each u_α is an ultrafilter of B. Since $f(X_0) \subseteq Y_0$, we obtain that each v_α is an ultrafilter of B'. Also, $u_\alpha \times u_\beta \subseteq (C_{\underline{X}})^{\#}$ for all $\alpha, \beta \in A$, i.e. $u_\alpha R_1' u_\beta$ for all $\alpha, \beta \in A$. Since $t_{\underline{X}}$ is a **PCS**-isomorphism, we get that $x_\alpha R_1 x_\beta$ for all $\alpha, \beta \in A$. Then $f(x_\alpha) R_2 f(x_\beta)$ for all $\alpha, \beta \in A$, and thus $t_{\underline{Y}}(f(x_\alpha)) R_2' t_{\underline{Y}}(f(x_\beta))$ for all $\alpha, \beta \in A$. This means that $v_a \times v_\beta \subseteq (C_{\underline{Y}})^{\#}$ for all $\alpha, \beta \in A$, and, hence, $\{v_\alpha \mid \alpha \in A\}$ is a clique in $(B', (C_{\underline{Y}})^{\#})$. Thus Γ' is a clan in $(B', (C_{\underline{Y}})^{\#})$, i.e. $\Gamma' \in Y'$. Then there exists $y \in Y$ such that $\Gamma' = t_{\underline{Y}}(y)$, i.e. $\Gamma' = \sigma_y^{B'}$. Now we show that $y = f(x)$ exactly as in the paragraph immediately after (26). So, (28) is proved. Analogously we get that $\sigma_{g(x)}^{B'} = \{\mathrm{cl}_Y(G') \mid G' \in CO(Y_0), g(x_\alpha) \in G' \text{ for some } \alpha \in A\}$. Let $g' = G^a(G^t(g))$. Then Theorem 3.4 implies that $g'(\sigma_x^B) = \sigma_{g(x)}^{B'}$. Since $f|_{X_0} = g|_{X_0}$, we get that $f'(\sigma_x^B) = g'(\sigma_x^B)$. Then $f(x) = g(x)$. So, $f = g$. □

Corollary 3.9. *The category* **PCS** *is equivalent to the category* **SAS** *of all Stone adjacency spaces and all continuous maps $f : (X_0, R) \longrightarrow (X_0', R')$ between them such that, for every $x, y \in X_0$, xRy implies $f(x)R'f(y)$.*

Proof. Let $F^t : \mathbf{PCS} \longrightarrow \mathbf{SAS}$ be the functor defined by

$$F^t(X, X_0, R) = (X_0, R)$$

on the objects of the category **PCS**, and by

$$F^t(f) = f|_{X_0}$$

for every $f \in \mathbf{PCS}((X, X_0, R), (Y, Y_0, R'))$. Clearly, F^t is a functor. We will show that it is full, faithful and isomorphism-dense, i.e. that it is an equivalence functor.

Let $(X_0, R) \in |\mathbf{SAS}|$. Set $\underline{B} = (CO(X_0), C_R)$ and $\underline{X} = G^a(\underline{B})$. According to Theorem 3.4, $\underline{X} \in |\mathbf{PCS}|$. Let $\underline{X} = (X', X_0', R')$ and $B = CO(X_0)$. Then, by the Stone duality theorem, $f : X_0 \longrightarrow X_0'$, $x \mapsto u_x$, is a homeomorphism (recall that $u_x = \{P \in B \mid x \in P\}$). We will show that for every $x, y \in X_0$, $xRy \iff f(x)R'f(y)$. Indeed, let $x, y \in X_0$ and xRy. We have that $f(x)R'f(y) \iff u_x R' u_y \iff u_x \times u_y \subseteq C_R \iff [(\forall P \in u_x)(\forall Q \in u_y)(PC_RQ)] \iff ([(x \in P \in B) \wedge (y \in Q \in B)] \Rightarrow (R \cap (P \times Q) \neq \emptyset))$. Now, since xRy, we obtain that $f(x)R'f(y)$. Conversely, let $x, y \in X_0$ and $f(x)R'f(y)$. Suppose that $x(-R)y$. Since R is a closed relation, there exist $P, Q \in B$ such that $x \in P$, $y \in Q$ and $R \cap (P \times Q) = \emptyset$. We get that $f(x)(-R')f(y)$, a contradiction. Hence, xRy. Therefore, f is a **SAS**-isomorphism

between (X_0, R) and (X_0', R'). Since $F^t(\underline{X}) = F^t(X', X_0', R') = (X_0', R')$, we obtain that F^t is isomorphism-dense.

Clearly, Corollary 3.8 implies that F^t is a faithful functor.

Let $\underline{X} = (X, X_0, R_0)$, $\underline{Y} = (Y, Y_0, R_1)$ and $\underline{X}, \underline{Y} \in |\mathbf{PCS}|$. We will show that the restriction of F^t to the hom-set $\mathbf{PCS}(\underline{X}, \underline{Y})$ is a surjection (i.e., the correspondence $\mathbf{PCS}(\underline{X}, \underline{Y}) \longrightarrow \mathbf{SAS}(F^t(\underline{X}), F^t(\underline{Y}))$, $f \mapsto f|_{X_0}$, is a surjection). Let $g \in \mathbf{SAS}((X_0, R_0), (Y_0, R_1))$ and $\varphi = S^t(g) : CO(Y_0) \longrightarrow CO(X_0)$, $G \mapsto g^{-1}(G)$, where $S^t : \mathbf{Stone} \longrightarrow \mathbf{Bool}$ is the Stone duality functor. In fact,

$$\varphi \in \mathbf{PCA}((CO(Y_0), C_{R_1}), (CO(X_0), C_{R_0})).$$

Indeed, let $G_1, G_2 \in CO(Y_0)$ and $\varphi(G_1) C_{R_0} \varphi(G_2)$. Then there exist $x \in \varphi(G_1)$ and $y \in \varphi(G_2)$ such that $x R_0 y$. We get that $x \in g^{-1}(G_1)$, $y \in g^{-1}(G_2)$ and $x R_0 y$. Thus $g(x) \in G_1$, $g(y) \in g(G_2)$ and $g(x) R_1 g(y)$. Hence, $G_1 C_{R_1} G_2$. So, φ is a **PCA**-morphism. Then Theorem 3.4 implies that

$$G^a(\varphi) \in \mathbf{PCS}(G^a(CO(X_0), C_{R_0}), G^a(CO(Y_0), C_{R_1})).$$

Let $G^a(CO(X_0), C_{R_0}) = (X', X_0', R_0')$ and $G^a((CO(Y_0), C_{R_1}) = (Y', Y_0', R_1')$. Then $X_0' = S(S^t(X_0))$ and $Y_0' = S(S^t(Y_0))$, where $S : \mathbf{Bool} \longrightarrow \mathbf{Stone}$ is the Stone duality functor. Thus $f_{X_0} : X_0 \longrightarrow X_0'$, $x \mapsto u_x$, and $f_{Y_0} : Y_0 \longrightarrow Y_0'$, $y \mapsto u_y$, are homeomorphisms and $f_{Y_0} \circ g = S(S^t(g)) \circ f_{X_0}$. Let $g' = S(S^t(g))$. Then we get that $u_{g(x)} = g'(u_x)$, for every $x \in X_0$. We will show that $g' = (G^a(\varphi))|_{X_0'}$. Indeed, for every $x \in X_0$, we have that $(G^a(\varphi))(u_x) = \varphi^{-1}(u_x) = \{G \in CO(Y_0) \mid \varphi(G) \in u_x\} = \{G \in CO(Y_0) \mid x \in \varphi(G)\} = \{G \in CO(Y_0) \mid x \in g^{-1}(G)\} = \{G \in CO(Y_0) \mid g(x) \in G\} = u_{g(x)} = g'(u_x)$. Hence, $F^t(G^a(\varphi)) = g'$. Since $(CO(X_0), C_{R_0})$ and $(RC(X, X_0), C_{\underline{X}})$ are **PCA**-isomorphic and, analogously, $(CO(Y_0), C_{R_1})$ and $(RC(Y, Y_0), C_{\underline{Y}})$ are **PCA**-isomorphic, we get that $G^a(CO(X_0), C_{R_0})$ is **PCS**-isomorphic to $G^a(G^t(\underline{X}))$ and, analogously, $G^a(CO(Y_0), C_{R_1})$ is **PCS**-isomorphic to $G^a(G^t(\underline{Y}))$. Now, using Theorem 3.4, we obtain that there exists $g_1 \in \mathbf{PCS}(\underline{X}, \underline{Y})$ such that $F^t(g_1) = g$. Thus, the functor F^t is full.

All this shows that $F^t : \mathbf{PCS} \longrightarrow \mathbf{SAS}$ is an equivalence functor. □

Corollary 3.10. ([3]) *The categories* **PCA** *and* **SAS** *are dually equivalent.*

Proof. Clearly, it follows from Theorem 3.4 and Corollary 3.9. □

Remark 3.11. It is not difficult to see that a direct proof of Corollary 3.10 can be easily obtained using Theorem 2.11.

Note that Corollary 3.10 is formulated in [3] using the relation *non-tangential inclusion* (defined here in (1); see also Remark 2.3) instead of the precontact relation; the relation non-tangential inclusion is called there *subordination*.

Corollary 3.12. *For every Stone adjacency space (X_0, R) there exists a unique (up to homeomorphism) topological space X such that the triple (X, X_0, R) is a 2-precontact space (and, thus, X is a compact semiregular T_0-space).*

Proof. By Corollary 3.9, there exists an equivalence functor $F^s : \mathbf{SAS} \longrightarrow \mathbf{PCS}$ such that the compositions $F^t \circ F^s$ and $F^s \circ F^t$ are naturally isomorphic to the corresponding identity functors. Let $(X_0, R) \in |\mathbf{SAS}|$. Since F^t is isomorphism-dense, there exists $(X', X'_0, R') \in |\mathbf{PCS}|$ such that $F^t(X', X'_0, R')$ is **SAS**-isomorphic to (X_0, R). Then, clearly, there exists $\underline{X} = (X, X_0, R) \in |\mathbf{PCS}|$. Suppose that there exists $\underline{X}_1 = (X_1, X_0, R) \in |\mathbf{PCS}|$. Then $F^s(F^t(\underline{X})) = F^s(F^t(\underline{X}_1))$ and thus $\underline{X}$ is **PCS**-isomorphic to $\underline{X}_1$. This implies that X is homeomorphic to X_1. Note that, by Definition 2.18 and Corollary 2.23, X is a compact semiregular T_0-space. □

Corollary 3.13. (A Duality Theorem for Contact Algebras) *The full subcategory* **CA** *of the category* **PCA** *whose objects are all contact algebras is dually equivalent to the category* **CS** *of all 2-contact spaces and all continuous maps* $f : (X, X_0) \longrightarrow (X', X'_0)$ *between them such that* $f(X_0) \subseteq X'_0$.

Proof. Let $\mathbf{S}''$ be the full subcategory of the category **PCS** whose objects are all 2-precontact spaces (X, X_0, R) for which R is a reflexive and symmetric relation. Then the categories S'' and **CS** are isomorphic. Indeed, let $F^c : \mathbf{S}'' \longrightarrow \mathbf{CS}$ be defined by $F^c(X, X_0, R) = (X, X_0)$ on the objects of the category $\mathbf{S}''$, and by $F^c(f) = f$ on the morphisms of the the category $\mathbf{S}''$. Then, using [16, Proposition 7.7], we get that F^c is well-defined and, obviously, it is a functor. By [16, Lemma 7.5], for every $(X, X_0) \in |\mathbf{CS}|$ there exists a unique relation $R_{(X,X_0)}$ on X_0 such that $(X, X_0, R) \in |\mathbf{S}''|$; as it is shown in [16, Lemma 7.5], the relation $R_{(X,X_0)}$ is defined as follows: for every $x, y \in X_0$,

$$(29)\quad xR_{(X,X_0)}y \iff ((\forall F \in u_x)(\forall G \in u_y)(\mathrm{cl}_X(F) \cap \mathrm{cl}_X(G) \neq \emptyset)),$$

where $u_x = \{A \in CO(X_0) \mid x \in A\}$ and analogously for u_y (see (7)). Now we set $F^d(X, X_0) = (X, X_0, R_{(X,X_0)})$, for every $(X, X_0) \in |\mathbf{CS}|$. Also, for every $f \in \mathbf{CS}((X, X_0), (Y, Y_0))$, we set $F^d(f) = f$. Then $f \in \mathbf{S}''(F^d(X, X_0), F^d(Y, Y_0))$. Indeed, set $R = R_{(X,X_0)}$ and $R' = R_{(Y,Y_0)}$. Let $x, y \in X_0$ and xRy. Then $f(x), f(y) \in Y_0$ (since $f(X_0) \subseteq Y_0$). Suppose that $f(x)(-R')f(y)$. Then there exist $F' \in u_{f(x)}$ and $G' \in u_{f(y)}$ such that $\mathrm{cl}_Y(F') \cap \mathrm{cl}_Y(G') = \emptyset$. Let $F = X_0 \cap f^{-1}(F')$ and $G = X_0 \cap f^{-1}(G')$. Then $F \in u_x$ and $G \in u_y$. Thus there exists $z \in \mathrm{cl}_X(F) \cap \mathrm{cl}_X(G)$. Since f is a continuous function, we get that $f(z) \in f(\mathrm{cl}_X(F)) \subseteq \mathrm{cl}_Y(f(F)) \subseteq \mathrm{cl}_Y(F')$ and, analogously, $f(z) \in \mathrm{cl}_Y(G')$. Therefore, $\mathrm{cl}_Y(F') \cap \mathrm{cl}_Y(G') \neq \emptyset$, a contradiction. Hence $f(x)R'f(y)$. So, $f \in \mathbf{S}''(F^d(X, X_0), F^d(Y, Y_0))$. All this shows that we have defined a functor $F^d : \mathbf{CS} \longrightarrow \mathbf{S}''$. Clearly, $F^c \circ F^d = Id_{\mathbf{CS}}$ and $F^d \circ F^c = Id_{\mathbf{S}''}$. Hence the categories $\mathbf{S}''$ and **CS** are isomorphic.

We will now show that $G^a(\mathbf{CA}) \subseteq \mathbf{S}''$ and $G^t(\mathbf{S}'') \subseteq \mathbf{CA}$. The first inclusion follows from [16, Lemma 3.5(d,e)], and the second one follows from Proposition 2.5(b) since, for every $\underline{X} = (X, X_0, R) \in |\mathbf{PCS}|$, the precontact algebras $(RC(X, X_0), C_{\underline{X}})$ and $(CO(X_0), C_R)$ are isomorphic. Now, applying Theorem 3.4, we obtain that the categories **CA** and **CS** are dually equivalent. □

Corollary 3.14. *Let* $f \in \mathbf{CS}((X, X_0), (Y, Y_0))$. *Then, for every* $H \in RC(Y, Y_0)$,

$$(30)\quad f^{-1}(H) = \mathrm{cl}_X(X_0 \cap f^{-1}(H)).$$

Proof. It follows from Corollary 3.13, Corollary 3.5 and the obvious fact that that $X_0 \cap f^{-1}(H) = X_0 \cap f^{-1}(H \cap Y_0)$ (because $f(X_0) \subseteq Y_0$). □

The next corollary follows immediately from Corollary 3.10 and Proposition 2.5:

Corollary 3.15. ([3]) *The category* **CA** *is dually equivalent to the full subcategory* **CSAS** *of the category* **SAS** *whose objects are all Stone adjacency spaces* (X, R) *such that* R *is a reflexive and symmetric relation.*

Let us derive it from Corollary 3.13 as well. Indeed, by the proof of Corollary 3.13, the categories $\mathbf{S}''$ and **CA** are dually equivalent; by Corollary 3.9, the functor $F^t : \mathbf{PCS} \longrightarrow \mathbf{SAS}$ is an equivalence; obviously, $F^t(\mathbf{S}'') = \mathbf{CSAS}$ and hence the categories $\mathbf{S}''$ and **CSAS** are equivalent; therefore, the categories **CSAS** and **CA** are dually equivalent.

Corollary 3.16. (A Duality Theorem for Complete Contact Algebras) *The full subcategory* **CCA** *of the category* **PCA** *whose objects are all complete contact algebras is dually equivalent to the category* **CSRS** *of all C-semiregular spaces and all continuous maps between them which preserve u-points.*

Proof. Let $\mathbf{S}'''$ be the full subcategory of the category **PCS** whose objects are all 2-precontact spaces (X, X_0, R) for which R is a reflexive and symmetric relation and X_0 is extremally disconnected. Then the categories S''' and **CSRS** are isomorphic. Indeed, let $F^e : \mathbf{S}''' \longrightarrow \mathbf{CSRS}$ be defined by $F^e(X, X_0, R) = X$ on the objects of the category $\mathbf{S}'''$, and by $F^e(f) = f$ on the morphisms of the the category $\mathbf{S}'''$. Then, by [16, Proposition 7.7] and Lemma 2.33, F^e is well-defined on the objects of the category S'''. Let $\underline{X} = (X, X_0, R) \in |\mathbf{S}'''|$, $\underline{Y} = (Y, Y_0, R') \in |\mathbf{S}'''|$ and $f \in \mathbf{S}'''(\underline{X}, \underline{Y})$. We will show that $F^e(f) \in \mathbf{CSRS}(F^e(\underline{X}), F^e(\underline{Y}))$. We have, by [16, Proposition 7.7], Lemma 2.33 and Theorem 2.32, that $X_0 = \{x \in X \mid x \text{ is an u-point of } X\}$ and $Y_0 = \{y \in Y \mid y \text{ is an u-point of } Y\}$. Since f is a continuous map and $f(X_0) \subseteq Y_0$, we get that f is a **CSRS**-morphism. So, F^e is well-defined. Obviously, F^e is a functor. Let now $X \in |\mathbf{CSRS}|$. We set $F^f(X) = (X, X_0, R_{(X,X_0)})$, where $X_0 = \{x \in X \mid x \text{ is an u-point of } X\}$ and the relation $R_{(X,X_0)}$ on X_0 is defined by (29). Then, using Theorem 2.32 and the proof of Corollary 3.13, we get that $F^f(X) \in |\mathbf{S}'''|$. Further, for every $f \in \mathbf{CSRS}(X, Y)$, we set $F^f(f) = f$. Let $F^f(X) = (X, X_0, R_{(X,X_0)})$ and $F^f(Y) = (Y, Y_0, R_{(Y,Y_0)})$. Then, clearly, $f(X_0) \subseteq Y_0$ and since, by Theorem 2.32, (X, X_0) and (Y, Y_0) are 2-contact spaces, we obtain that $f \in \mathbf{CS}((X, X_0), (Y, Y_0))$; now, by Corollary 3.13, $f = F^d(f)$ and we get that $f \in \mathbf{S}''(F^d(X, X_0), F^d(Y, Y_0))$; this implies that $f \in \mathbf{S}'''(F^f(X), F^f(Y))$. All this shows that we have defined a functor $F^f : \mathbf{CSRS} \longrightarrow \mathbf{S}'''$. Clearly, $F^e \circ F^f = Id_{\mathbf{CSRS}}$ and $F^f \circ F^e = Id_{\mathbf{S}'''}$. Hence the categories $\mathbf{S}'''$ and **CSRS** are isomorphic.

We will now show that $G^a(\mathbf{CCA}) \subseteq \mathbf{S}'''$ and $G^t(\mathbf{S}''') \subseteq \mathbf{CCA}$. By Corollary 3.13, we have that $G^a(\mathbf{CCA}) \subseteq \mathbf{S}''$ and then Theorem 2.22(a) together with Lemma 2.14 imply that even $G^a(\mathbf{CCA}) \subseteq \mathbf{S}'''$. Further, by Corollary 3.13, $G^t(\mathbf{S}'') \subseteq \mathbf{CA}$. Since, for every $\underline{X} = (X, X_0, R) \in |\mathbf{S}'''|$, we have that $CO(X_0) = RC(X_0)$ and thus $RC(X, X_0) = RC(X)$, we obtain that $G^t(\mathbf{S}''') \subseteq \mathbf{CCA}$. Now, applying Theorem 3.4, we obtain that the categories **CCA** and **CSRS** are dually equivalent. □

We are now going to present Corollary 3.13 in a form similar to that of Corollary 3.16. For doing this we need to recall two definitions from [23]. We first introduce a new notion.

Definition 3.17. Let X be a topological space and B be a Boolean subalgebra of the Boolean algebra $RC(X)$. Then the pair (X, B) is called a *mereotopological pair*.

Definition 3.18. ([23]) (a) A mereotopological pair (X, B) is called a *mereotopological space* if B is a closed base for X. We say that (X, B) is a mereotopological T_0-space if (X, B) is a mereotopological space and X is a T_0-space.

(b) A mereotopological space (X, B) is said to be *mereocompact* if for every clan Γ of $(B, C_X \cap B^2)$ there exists a point x of X such that $\Gamma = \sigma_x^B$ (see (9) for σ_x^B).

Remark 3.19. (a) Obviously, if (X, B) is a mereotopological space then X is semiregular.

(b) The definition of mereocompactness given here is one of the several equivalent expressions of the definition of mereocompactness given in [23]. We will use only the definition given here.

(c) Clearly, a space X is C-semiregular (see Definition 2.29) iff $(X, RC(X))$ is a mereocompact T_0-space. So that, the notion of mereocompactness is an analogue of the notion of C-semiregular space.

Having in mind the definition of an u-point of a topological space (see Definition 2.31), we will now introduce the more general notion of an *u-point of a mereotopological pair*.

Definition 3.20. Let (X, B) be a mereotopological pair and $x \in X$. Then the point x is said to be an *u-point of the mereotopological pair* (X, B) if, for every $F, G \in B$, $x \in F \cap G$ implies that $x \in \mathrm{cl}_X(\mathrm{int}_X(F \cap G))$.

Obviously, a point x of a topological space X is an u-point of X iff it is an u-point of the mereotopological pair $(X, RC(X))$. Also, if X is a topological space, then every point of X is an u-point of the mereotopological pair $(X, CO(X))$.

We will now generalize some assertions from [16] concerning u-points.

Proposition 3.21. *Let (X, X_0) be a topological pair, (X, B) be a mereotopological pair (resp., a mereotopological space) and $B_0 = \{F \cap X_0 \mid F \in B\}$. Then (X_0, B_0) is a mereotopological pair (resp., a mereotopological space); also, for every $x \in X_0$, we have that x is an u-point of (X_0, B_0) iff x is an u-point of (X, B).*

Proof. Using Lemma 2.12, we get that (X_0, B_0) is a mereotopological pair (resp., a mereotopological space). Note that, by Lemma 2.12, for every $H \in B$ there exist $H_0 \in B_0$ such that $H = \mathrm{cl}_X(H_0)$ and, also, for every $H_0 \in B_0$, $\mathrm{cl}_X(H_0) \in B$. Let $x \in X_0$ be an u-point of (X_0, B_0), $F, G \in B_0$ and $x \in \mathrm{cl}_X(F) \cap \mathrm{cl}_X(G)$. Then $x \in F \cap G$ and hence $x \in \mathrm{cl}_{X_0}(\mathrm{int}_{X_0}(F \cap G))$. Since, by Lemma 2.12, $\mathrm{cl}_X(cl_{X_0}(\mathrm{int}_{X_0}(F \cap G))) = \mathrm{cl}_X(\mathrm{int}_X(\mathrm{cl}_X(F) \cap \mathrm{cl}_X(G)))$, we get that x is an u-point of (X, B). Conversely, let $x \in X_0$ be an u-point of (X, B). Let $F, G \in B_0$ and $x \in F \cap G$. Then $x \in$

$\mathrm{cl}_X(F) \cap \mathrm{cl}_X(G)$ and thus $x \in \mathrm{cl}_X(\mathrm{int}_X(\mathrm{cl}_X(F) \cap \mathrm{cl}_X(G)))$. Since, by Lemma 2.12, $X \cap \mathrm{cl}_X(\mathrm{int}_X(\mathrm{cl}_X(F) \cap \mathrm{cl}_X(G))) = \mathrm{cl}_{X_0}(\mathrm{int}_{X_0}(F \cap G))$, we get that x is an u-point of (X_0, B_0). □

Proposition 3.22. *Let (X, B) be a topological pair and $x \in X$. Then x is an u-point of (X, B) iff σ_x^B is an ultrafilter of the Boolean algebra B.*

Proof. Since, by (20), σ_x^B is a clan of $(B, C_X \cap B^2)$, we obtain that σ_x^B is a grill of B. Hence: (σ_x^B is an ultrafilter of B) $\iff$ $[(\forall F, G \in \sigma_x^B)(F.G \in \sigma_x^B)]$ $\iff$ (x is an u-point of (X, B)). (Note that $F.G = cl_X(\mathrm{int}_X(F \cap G))$.) □

Theorem 3.23. *For every mereocompact T_0-space (X, B), the set*

$$u(X, B) = \{x \in X \mid x \text{ is an u-point of } (X, B)\}$$

endowed with its subspace topology is a dense zero-dimensional compact Hausdorff subspace of X and is the unique dense zero-dimensional compact Hausdorff subspace of X such that $RC(X, u(X, B)) = B$. Also, the pair $(X, u(X, B))$ is a 2-contact space.

Proof. Set $X_0 = u(X, B)$. We have that $\underline{B} = (B, C_X \cap B^2)$ is a contact algebra. Hence, by Definition 2.25(c), $\underline{X}^c(\underline{B})$ is a 2-contact space. Let $\underline{X}^c(\underline{B}) = (X', X_0')$. Then $X' = Clans(\underline{B})$ and $X_0' = Ult(B)$. Since X is a T_0-space and B is a closed base for X, arguing as in the paragraph after (22) and using (20), we obtain that the map

$$t_X^c : X \longrightarrow X', \quad x \mapsto \sigma_x^B,$$

is an injection. Also, the fact that (X, B) is mereocompact implies that t_X^c is a surjection. Now, arguing as in the paragraph immediately after (23), we get that t_X^c is a homeomorphism and

(31) $\forall F \in B,\ t_X^c(F) = g_{\underline{B}}(F)$.

Using Proposition 3.22, we obtain that $t_X^c(X_0) = X_0'$. All this shows that (X, X_0) is a 2-contact space. Hence, X_0 is a Stone space and X_0 is dense in X. Using (31) and Theorem 2.22(a) (or [16, Theorem 7.9(a)]), we get that $B = RC(X, X_0) = \{\mathrm{cl}_X(P) \mid P \in CO(X_0)\}$.

For proving the uniqueness of X_0, let X_0^1 be a dense Stone subspace of X such that $B = \{\mathrm{cl}_X(P) \mid P \in CO(X_0^1)\}$. We will show that $X_0^1 = X_0$. Set $B_0 = CO(X_0^1)$. Then $B_0 = \{F \cap X_0^1 \mid F \in B\}$ (by Lemma 2.12) and (X_0^1, B_0) is a mereotopological space. Obviously, every point of X_0^1 is an u-point of (X_0^1, B_0). Then Proposition 3.21 implies that every point of X_0^1 is an u-point of (X, B). Thus $X_0^1 \subseteq X_0$. Obviously, X_0^1 is a dense subspace of X_0. Since X_0^1 is compact and X_0 is Hausdorff, we get that $X_0^1 = X_0$. □

Note that Theorem 3.23 and Proposition 2.26 imply the following result from [23]: if (X, B) is a mereocompact T_0-space then X is compact. (In fact, a stronger result is proved in [23]: every mereocompact space is compact.)

Lemma 3.24. *If (X, X_0) is a 2-contact space, then $(X, RC(X, X_0))$ is a mereocompact T_0-space.*

Proof. Clearly, the map $e_{X_0,X} : (CO(X_0), \delta_{(X,X_0)}) \longrightarrow (RC(X, X_0), \quad F \mapsto \mathrm{cl}_X(F)$, is a CA-isomorphism (use Lemma 2.12) and for every $x \in X$, $e_{X_0,X}(\Gamma_{x,X_0}) = \sigma^B_x$. Then the axioms (CS1), (CS3) and (CS4) from Definition 2.25 imply that $(X, RC(X, X_0))$ is a mereocompact T_0-space. □

Definition 3.25. Let us denote by **MCS** the category whose objects are all mereocompact T_0-spaces and whose morphisms are all continuous maps between mereocompact T_0-spaces which preserve the corresponding u-points (i.e.,

$$f \in \mathbf{MCS}((X, A), (Y, B))$$

iff $f : X \longrightarrow Y$ is a continuous map and, for every u-point x of (X, A), $f(x)$ is an u-point of (Y, B)).

Obviously, **MCS** is indeed a category.

Theorem 3.26. (A Duality Theorem for Contact Algebras) *The categories* **CA** *and* **MCS** *are dually equivalent.*

Proof. Having in mind Corollary 3.13, it is enough to show that the categories **CS** and **MCS** are isomorphic. Let $F^g : \mathbf{CS} \longrightarrow \mathbf{MCS}$ be defined by $F^g(X, X_0) = (X, RC(X, X_0))$ on the objects of the category **CS**, and by $F^g(f) = f$ on the morphisms of the category **CS**. Then Lemma 3.24 shows that F^g is well defined on the objects of the category **CS**. Let $f \in \mathbf{CS}((X, X_0), (Y, Y_0))$. Then $f : X \longrightarrow Y$ is a continuous function and $f(X_0) \subseteq Y_0$. Since (X, X_0) is a 2-contact space, we get that X_0 is a dense Stone subspace of X. Since $(X, RC(X, X_0))$ is a mereocompact T_0-space and $RC(X, X_0) = \{\mathrm{cl}_X(P) \mid P \in CO(X_0)\}$, Theorem 3.23 implies that $X_0 = \{x \in X \mid x \text{ is an u-point of } (X, RC(X, X_0))\}$. Analogously, we get that $Y_0 = \{y \in Y \mid y \text{ is an u-point of } (Y, RC(Y, Y_0))\}$. Since $f(X_0) \subseteq Y_0$, we get that f preserves the u-points. Hence, $F^g(f) \in \mathbf{MCS}(F^g(X, X_0), F^g(Y, Y_0))$. So, F^g is well defined on the morphisms as well. Obviously, F^g is a functor.

Let $F^h : \mathbf{MCS} \longrightarrow \mathbf{CS}$ be defined by $F^h(X, B) = (X, u(X, B))$ (see Theorem 3.23 for the notation $u(X, B)$) on the objects of the category **MCS**, and by $F^h(f) = f$ on the morphisms of the category **MCS**. Then Theorem 3.23 implies that F^h is well defined. Obviously, F^h is a functor.

Using once more Theorem 3.23, we get that $F^g \circ F^h = Id_{\mathbf{MCS}}$ and $F^h \circ F^g = Id_{\mathbf{CS}}$. Hence, the categories **CS** and **MCS** are isomorphic. □

Finally, we will show how our results imply the duality for contact algebras described in [23].

Definition 3.27. ([23]) Let **GMCS** be the category whose objects are all mereocompact T_0-spaces and whose morphisms are defined as follows:

$$f \in \mathbf{GMCS}((X, A), (Y, B))$$

iff $f : X \longrightarrow Y$ is a function such that the function

$$\psi_f : B \longrightarrow A, \quad F \mapsto f^{-1}(F),$$

is well defined and is a Boolean homomorphism.

Corollary 3.28. ([23]) *The categories* **CA** *and* **GMCS** *are dually equivalent.*

Proof. We will derive this result from Corollary 3.13 showing that the categories **CS** and **GMCS** are isomorphic. Let $F^i : \mathbf{CS} \longrightarrow \mathbf{GMCS}$ be defined by $F^i(X, X_0) = (X, RC(X, X_0))$ on the objects of the category **CS**, and by $F^i(f) = f$ on the morphisms of the category **CS**. Then Lemma 3.24 shows that F^i is well defined on the objects of the category **CS**. Let $f \in \mathbf{CS}((X, X_0), (Y, Y_0))$. Then $f : X \longrightarrow Y$ is a continuous function and $f(X_0) \subseteq Y_0$. Now Corollary 3.14, (13) and (14) show that the function $\psi_f : RC(Y, Y_0) \longrightarrow RC(X, X_0),\ H \mapsto f^{-1}(H)$, is a Boolean homomorphism. Indeed, we have that $H \in RC(Y, Y_0) \iff H = \mathrm{cl}_Y(F)$, where $F \in CO(Y_0)$; thus, for every $H \in RC(Y, Y_0)$, $H \cap Y_0 \in CO(Y_0)$. Therefore, $f^{-1}(H) = \mathrm{cl}_X(X_0 \cap f^{-1}(H \cap Y_0))$. Then, using formula (13) and arguing as in the paragraph after it, we get that ψ_f is a Boolean homomorphism. Hence F^i is well defined on the morphisms of the category **CS** as well. Obviously, F^i is a functor.

Let $F^j : \mathbf{GMCS} \longrightarrow \mathbf{CS}$ be defined by $F^j(X, B) = (X, u(X, B))$ (see Theorem 3.23 for the notation $u(X, B)$) on the objects of the category **GMCS**, and by $F^j(f) = f$ on the morphisms of the category **GMCS**. Then Theorem 3.23 implies that F^j is well defined on the objects of the category **GMCS**. If $f \in \mathbf{GMCS}((X, A), (Y, B))$ then f is a continuous map because B is a closed base of Y and $f^{-1}(B) \subseteq A$. Let $F^j(X, A) = (X, X_0)$ and $F^j(Y, B) = (Y, Y_0)$. We have to show that $f(X_0) \subseteq Y_0$, i.e. that if x is an u-point of (X, A) then $f(x)$ is an u-point of (Y, B). So, let x be an u-point of (X, A), $F, G \in B$ and $f(x) \in F \cap G$. Then $x \in f^{-1}(F) \cap f^{-1}(G)$ and $f^{-1}(F), f^{-1}(G) \in A$. Hence $x \in f^{-1}(F).f^{-1}(G) = \psi_f(F).\psi_f(G) = \psi_f(F.G) = f^{-1}(F.G)$. Therefore, $f(x) \in F.G$ and thus $f(x)$ is an u-point of (Y, B). So, F^j is well defined on the morphisms of the category **GMCS**. Obviously, F^j is a functor.

Using once more Theorem 3.23, we get that $F^i \circ F^j = Id_{\mathbf{GMCS}}$ and $F^j \circ F^i = Id_{\mathbf{CS}}$. Hence, the categories **CS** and **GMCS** are isomorphic. □

Since, by Theorem 2.22(a), a precontact algebra $\underline{B}$ is connected iff its canonical 2-precontact space $\underline{X}(\underline{B})$ is connected, all our results from this section which concern dualities have as corollaries duality theorems for the corresponding full subcategories of the corresponding categories of (pre)contact algebras having as objects all connected (pre)contact algebras. We will now formulate, as an example, the connected variant of Theorem 3.4 and will left to the reader the formulation of all other corollaries.

Theorem 3.29. *The full subcategory* **PCAC** *of the category* **PCA** *whose objects are all connected precontact algebras is dually equivalent to the full subcategory* **PCSC** *of the category* **PCS** *having as objects all connected 2-precontact spaces.*

References

[1] Adámek, J., Herrlich, H. and Strecker, G. E. *Abstract and Concrete Categories*,

[2] Alexandroff, P. S. and Ponomarev, V. I. On bicompact extensions of topological spaces, Vestn. Mosk. Univ. Ser. Mat. (1959), 93-108. (In Russian).

[3] Bezhanishvili, G., Bezhanishvili, N., Sourabh, S. and Venema, Y. Subordinations, closed relations, and compact Hausdorff spaces. (December 2014) pp. 1-21, available at http://www.phil.uu.nl/~bezhanishvili/Papers/stone_closed_relation_9.pdf

[4] Čech, E. *Topological Spaces.* Interscience, London, 1966.

[5] Comfort, W. and Negrepontis, S. *Chain Conditions in Topology*, Cambridge Univ. Press, Cambridge, 1982.

[6] de Vries, H. *Compact Spaces and Compactifications, an Algebraic Approach*, Van Gorcum, The Netherlands, 1962.

[7] Dimov, G. A generalization of De Vries' Duality Theorem, Applied Categorical Structures 17 (2009), 501-516.

[8] Dimov, G. Some generalizations of the Fedorchuk duality theorem - I, Topology Appl., 156 (2009), 728-746.

[9] Dimov, G. A de Vries-type duality theorem for the category of locally compact spaces and continuous maps - I, Acta Math. Hungarica, 129 (4) (2010), 314-349.

[10] Dimov, G. A de Vries-type duality theorem for the category of locally compact spaces and continuous maps - II, Acta Math. Hungarica, 130 (1) (2011), 50-77.

[11] Dimov, G. Some Generalizations of the Stone Duality Theorem, Publicationes Mathematicae Debrecen 80(3-4) (2012), 255-293.

[12] Dimov, G. and E. Ivanova Yet another duality theorem for locally compact spaces, Houston Journal of Mathematics (2016) (to appear).

[13] Dimov, G. and Vakarelov, D. Contact Algebras and Region-based Theory of Space: A Proximity Approach - I. Fundamenta Informaticae 74(2-3) (2006), 209-249.

[14] Dimov, G. and Vakarelov, D. Contact Algebras and Region-based Theory of Space: A Proximity Approach - II. Fundamenta Informaticae 74 (2-3) (2006), 251-282.

[15] Dimov, G. and Vakarelov, D. Topological Representation of Precontact Algebras. In: *Relation Methods in Computer Science*, W. MacCaull, M. Winter, I. Duentsch (Eds.), Lecture Notes in Computer Science, 3929 (2006), 1-16, Springer-Verlag Berlin Heidelberg.

[16] Dimov, G. and Vakarelov, D. Topological Representation of Precontact Algebras and a Connected Version of the Stone Duality Theorem – I.

arXiv:1508.02220v3, 1-44 (a revised version of this paper will appear in Topology Appl.).

[17] Düntsch, I. and Vakarelov, D. Region-based theory of discrete spaces: A proximity approach. In: Nadif, M., Napoli, A., SanJuan, E., and Sigayret, A. EDS, *Proceedings of Fourth International Conference Journées de l'informatique Messine*, 123-129, Metz, France, 2003. Journal version in: *Annals of Mathematics and Artificial Intelligence*, 49(1-4) (2007), 5-14.

[18] Düntsch, I. and Winter, M. A Representation theorem for Boolean Contact Algebras. Theoretical Computer Science (B), 347 (2005), 498-512.

[19] Efremovič, V. A. Infinitesimal spaces. DAN SSSR, 76 (1951), 341–343.

[20] Engelking, R. *General Topology*, PWN, Warszawa, 1977.

[21] Fedorchuk, V. V. Boolean δ-algebras and quasi-open mappings. Sibirsk. Mat. Ž. 14 (5) (1973), 1088–1099; English translation: Siberian Math. J. 14 (1973), 759-767 (1974).

[22] Galton, A. The mereotopology of discrete spaces. In: Freksa, C. and Mark, D.M. eds, *Spatial Information Theory, Proceedings of the International Conference COSIT'99,* Lecture Notes in Computer Science, 251–266, Springer-Verlag, 1999.

[23] Goldblatt, R. and Grice, M. Mereocompactness and duality for mereotopological spaces. In: Katalin Bimbo (Ed.), J. Michael Dunn on Information Based Logics, Springer, 2016 (to appear).
Available at http://homepages.ecs.vuw.ac.nz/∼rob/papers/mereo.pdf

[24] Koppelberg, S. *Handbook on Boolean Algebras, vol. 1: General Theory of Boolean Algebras*, North Holland, 1989.

[25] Naimpally, S. and Warrack, B. *Proximity Spaces.* Cambridge, London, 1970.

[26] Smirnov, J. M. On proximity spaces. Mat. Sb. 31 (1952), 543–574.

[27] Stone, M. H. The theory of representations for Boolean algebras. Trans. Amer. Math. Soc., 40 (1936), 37–111.

[28] Thron, W. Proximity structures and grills. *Math. Ann.*, 206 (1973), 35-62.

[29] Vakarelov, D., Dimov, G., Düntsch, I., Bennett, B. A proximity approach to some region-based theory of space. *Journal of applied non-classical logics*, 12 (3-4) (2002), 527-559.